




UNIVERSITY OF TORONTO



3 1761 00852744 2

UNIV. OF
TORONTO
LIBRARY





Digitized by the Internet Archive
in 2008 with funding from
Microsoft Corporation

ROYAL BOTANIC GARDENS, KEW.

H A N D - L I S T

OF

FERNS AND FERN ALLIES

CULTIVATED IN

THE ROYAL BOTANIC GARDENS.

(SECOND EDITION.)



L O N D O N :

SOLD AT THE ROYAL BOTANIC GARDENS, KEW.
PRINTED FOR HIS MAJESTY'S STATIONERY OFFICE,
BY DARLING & SON, LTD., 34-40, BACON STREET, E.

1906.

Price Fivepence.

86298
198
298
398
108

QK

73

K4A237

1906

PREFACE:

THE general purpose of the series of Hand-lists of the plants cultivated in the Royal Gardens which is in course of publication, and of which the present one is a part, is sufficiently indicated in the preface to that which enumerates the trees and shrubs.

The collection of Ferns, whether tropical or temperate, is perhaps, next to that of Palms, the most important feature of the cultivation under glass in the Royal Gardens.

A special interest attaches to it, because it has been used so much in the preparation of most of the books and papers dealing with ferns which have been published in this country during the last half century. The celebrated botanical artist, Francis Bauer, was resident draughtsman for 50 years (1790–1840) to the Royal Botanic Gardens at Kew, and the beautiful series of analytical drawings of ferns which he made were published by Sir William Hooker, after his death in 1842, with descriptive letter-press and additions, under the title of *Genera Filicum*. During the 25 years (1840–1865) for which Sir William Hooker was the director of the gardens, his attention was specially concentrated on ferns. Under the title of *Species Filicum* he published an elaborate work in five

volumes, in which all the known species of typical ferns are fully described, and their synonymy and geographical distribution worked out, with copious illustrations by Mr. Walter Fitch. Of this, the first volume appeared in 1846, and the last in 1864. He planned also a condensed hand-book of all the known ferns, to be called *Synopsis Filicum*, intended specially for the use of colonists and travellers. All the drawings for this had been made, but the letter-press was not far advanced when he died in the autumn of 1865. It was completed by Mr. J. G. Baker, F.R.S., now Keeper of the Herbarium, in 1868, and a second edition was brought out in 1874. Since that date many hundred new species have been discovered, and a volume of Hooker's *Icones Plantarum*, containing plates and descriptions of the more interesting of the novelties, was published in 1887. Mr. Baker has also published in the *Annals of Botany* (1891) a summary of new ferns discovered or described since 1874. The type specimens from which most of the descriptions and plates contained in this series of books have been made are deposited in the Kew Herbarium.

The living collection in the garden owes its completeness very largely to the zeal and assiduity with which the veteran pteridologist, Mr. John Smith, curator of the Royal Gardens from 1841-63, watched over it for more than 40 years.

In his privately printed *Records of Kew* (pp. 322, 323) he gives the following particulars of its origin and development:—"In the year 1822 I found the collection of ferns at Kew extremely poor, especially as regards tropical species, very many of those introduced in previous years

having been lost and very few new ones added. . . . The tender exotic species were more or less of them growing without any arrangement in different houses, and unnamed, their number amounting to about 40. In 1825 I arranged the tropical species in a group at the end of the then lean-to house . . . now included in the tropical fern house, the area they occupied being 6 ft. by 12 ft. These formed the nucleus of the now great collection. They were successively added to by importations of living plants, as also plants raised from spores obtained from herbarium specimens."

"The collection continued yearly to increase, and in 1846 [1845] I drew up a list of the collection, which was published in an appendix to the *Botanical Magazine*, for that year; the number then amounted to 400 [378] species. This in 1857 [1856] was followed by another list, entitled 'Catalogue of Cultivated Ferns,' in which 600 [504] species are enumerated."

The principal books published by Mr. Smith are his *Ferns, British and Foreign*, issued in 1866, which contains a classified list of all the species then known in cultivation, full directions for the cultivation of ferns of the different climatic types and by far the most complete history of their gradual introduction which has ever appeared in print, and his *Historia Filicum*, issued in 1875, which contains woodcuts of 220 types and gives a full exposition of his views on fern classification.

In 1868 the last published list was prepared by Mr. J. G. Baker. It enumerates 802 species and varieties of ferns and 48 of fern allies.

The present list comprises 1116 species and varieties of ferns and 97 of fern allies ; this is exclusive of British ferns, of which 586 varieties are in cultivation.

The collection falls into three great groups :—(i.) Tropical ; (ii.) Temperate ; and (iii.) Hardy Ferns.

The Tropical Ferns are cultivated in No. II. House.

The Temperate Ferns are cultivated in No. III. ; the Filmy Ferns have a separate House (No. II. A). constructed for the special treatment which they require. The fine collection of Temperate Tree Ferns is contained in the Temperate House.

The Hardy Ferns are arranged on and about a small rockery, $\frac{7}{E}$ of the Key Plan, and forming the south boundary of Lawn L.

The structural development of the buildings in which the collection is housed has kept pace with its growth. Their history may be given briefly.

The Tropical Fern House (No. II.) is a span-roofed house 129 feet long, 34 feet broad, and 15 feet high in the centre. It has a wide transect on the south side, 40 feet long, 33 feet wide, and 19 feet high.

In 1841 the site of this house was occupied by two lean-to houses used for miscellaneous collections of stove plants. They are enumerated in Dr. Lindley's Report to the Treasury (1838) as Nos. 2 and 3 ; the former stood to the west, and was 50 feet long ; the length of the other

was 60 feet. It was in No. 2 that, as stated above, John Smith first arranged the nucleus of the collection of Tropical Ferns.

In 1843, No. 2 was doubled and made into a span house.

In 1845, No. 3 was entirely taken down, and a new span-roofed house erected in its place. It joined on to No. 2, which was kept at a somewhat lower temperature, but opened into it through a partition. The centre was filled up with slate shelf staging, and the sides with stone shelves. No. 2 was used as a stove for rare palms and large tropical ferns. No. 3 was at first occupied by the Woburn collection of orchids presented by Her Majesty the Queen ; for this purpose it proved eventually unsuited. The orchids were gradually replaced by *Aroideæ* and ferns. Ultimately the latter obtained possession of the whole house. According to John Smith (Records, p. 334), "the centre of the house was a raised sloping rockery, amongst which the plants were growing in a natural state, forming fine bushy specimens."

A small house for tree ferns was built in 1861 ; this was connected with the principal house in 1868, and now forms the transept.

In 1879 the Tropical Fern House, which from the decay of the timbers had lapsed into an almost ruinous condition, was further severely damaged by the hail storm of August 3rd ; 1,152 panes of glass in it were broken. It was subsequently put into a thorough state of repair, but the collections received considerable damage.

In 1887 the west end of the house was set back a few feet ; the staging was partly re-arranged and the broad path through the transept was made.

The humid conditions necessary for the cultivation of tropical ferns are unfortunately not conducive to the preservation of the structures in which they are grown when they are built of wood. In 1889 it therefore again became necessary to renew the east wing of No. II. House. In order to secure greater durability it was decided to use iron for the principal framework, and wood only for the lights. The top ventilation was provided for by means of a lantern instead of by the older and more clumsy method of sliding sashes. At the same time the use of green glass in the reconstructed wing was abandoned. This was introduced at Kew in 1846 on the recommendation of the late Mr. Robert Hunt, F.R.S., who advocated its use on theoretical grounds which are now known to be mistaken. The resulting improvement in the growth of the plants, especially the species of *Adiantum* was very marked, and in 1892 a portion of the west wing was re-glazed with equally satisfactory results.

The temperate ferns are now collected in a house (No. III.), which was also built on the system of mixed iron and wood construction. It was erected in 1892 to take the place of a decayed and obsolete structure composed of two small greenhouses united together, Nos. 4 and 5 of Dr. Lindley's report. The former (the northern wing) was erected in 1803, and in 1840 contained New Holland and Cape plants ; the latter, which was "re-modelled" in 1825, was filled with succulents.

The reconstructed No. III. is a span-roofed house 60 feet long, 23 feet wide, and 13 feet high. It contains a broad central and two side stages, with paths on both sides.

The north wing of the former No. III. had been latterly devoted almost exclusively to filmy ferns after the Kew collection had been enriched by the gift of that formed by the late John Cooper Forster, which was presented to the Royal Gardens in 1888 by his widow.

The collection is now contained in the new Filmy Fern House (No. II. A.) erected on the north side of No. II. It is 50 feet long by 14 feet wide, with a central path and two cases running the full length of the house.

The hardy ferns occupy the rockery originally devoted to Alpine plants (before the construction of the Rock Garden in 1882), and it extends over part of the adjacent ground. A collection was established here in 1874, and the rockery itself was reconstructed in 1888.

The collection as a whole is now probably the richest in existence in garden forms of British species. This is due to the munificent bequest by W. C. Carbonell, Esq., of the extensive collection formed by him at Rhiw Castell, Usk, Monmouthshire. The whole was removed to Kew in 1887. It consisted of 4,261 specimens, many probably unique, besides some hundreds of seedlings.

The total number of well-marked species of ferns and the vascular Cryptogamia (fern allies) now mounts up to 3,500. Not more than one-third of these have been

brought into cultivation, and Kew is always glad to receive any additions, if possible in the form of well-established plants; if not, in the shape of dry spores, which can be collected and sent very easily by post in small packets.

The following table, which shows the percentage of the total number of ferns found in different parts of the world was drawn up by Mr. Baker in 1867; but he does not think that the discoveries of the last 20 years will have altered any of the figures materially :—

Arctic Zone	1 per cent.
Europe	4 ,,
Temperate Asia, including Himalayas	18 ,,
Temperate N. America	5 ,,
Temperate S. Africa	7 ,,
Australia and New Zealand	9 ,,
South temperate America	5 ,,
Tropical Africa	15 ,,
Tropical Asia	39 ,,
Tropical America...	42 ,,

The present hand list is divided into three parts :—

- (i.) Ferns proper (pp. 13–133);
- (ii.) Fern allies (pp. 135–143); and
- (iii.) as an appendix, a list of garden forms of British species (pp. 145–183).

With regard to (i.) ferns proper, the arrangement is alphabetical and synonyms have been only sparingly introduced, those included being chiefly such as are in

use in gardens and which are here referred to the correct name in each case. It has not been thought necessary, therefore, to supply an index. Those who wish to pursue the study of fern-nomenclature further may fall back on Hooker and Baker's *Synopsis Filicum*, upon which the present hand-list is substantially based.

The fern allies (ii.) have also been grouped alphabetically. It has not been thought necessary for the present purpose to separate them according to a botanical classification.

The cultivated and feral varieties of British ferns (iii.) have been enumerated in an appendix. Although the collection of them at Kew is very rich, they stand in a different position to the collection of recognised and well-determined species. Apart from their intrinsic beauty, which is often striking, they are of considerable scientific interest as showing the range of variation due to crossing and seminal reproduction. The amateurs and cultivators who have raised them have furnished them with Latin names, often cumbrous and fantastic, which have received no formal definition. They cannot, therefore, be fixed or quoted for any scientific purpose ; they have, in fact, the same relationship to the species from which they have originated as bedding pelargoniums bear to *Pelargonium zonale* or as the drumhead and other cabbages to *Brassica oleracea*.

PREFACE TO SECOND EDITION.

THE edition of the *Hand-List of Ferns and Fern-Allies cultivated in the Royal Gardens* which was issued in 1895 has for some time been exhausted. Since its publication some change has taken place in the composition of the Fern-collection. The present edition, the preparation of which for the press has been entrusted to Mr. C. H. Wright, A.L.S., Assistant in the Herbarium, is an attempt to enumerate the species and varieties now in cultivation at Kew. The species that have been acquired since 1895 are here included ; those that are not now in cultivation at Kew have been omitted. A few casual errors that had crept into the previous edition have been corrected. To facilitate the identification of the plants selected references have been given to publications in which reliable figures are to be found. In other respects the present edition follows the lines on which the edition of 1895 was prepared.

The table drawn up by Mr. J. G. Baker, F.R.S., in 1867, to show the percentage of the total number of ferns that had been collected up to that time in different parts of the world, which is included in the preface to the 1895 edition of this hand-list, is found, in the light of our present knowledge of the groups dealt with, to require revision. It will not be possible, until some general list, such as Carl Christensen's admirable *Index Filicum* (now in course of publication) is available,

to give absolute percentages. But the subjoined table, which may be taken as approximately accurate, will, it is hoped, be useful both in itself and for comparison with Mr. Baker's table, which remains of great historical interest :—

Arctic Zone4 per cent.
Europe	2·5	..
Temperate Asia, including the Himalayas above 8,000 ft.	...	25·5	"	
North Africa, Canary Is.8	"	
Temperate N. America	...	3·2	"	
Temperate S. Africa	...	4·5	"	
Australia and New Zealand	...	6·9	"	
Polynesia	10·8	"	
Temperate S. America	...	2·7	"	
Tropical Africa and Mascarene Is.	15·2	"		
Tropical Asia	...	34·8	"	
Tropical America	...	46·4	"	

D. P.

Kew. March, 1906.

I.—FERNS.

Aerophorus. *See* *Davallia.*

Acrostichum, L.

A. acuminatum, Hook. Brazil.

A. alienum, Sw. West Indies, Tropical America.

A. apodum, Kaulf. West Indies, Tropical America and Chile. Hook. & Grev. Ic. Fil. t. 99.

A. aureum, L. Tropical and sub-tropical regions.

A. auritum, Sw. Malaya, *Polybotrya aurita*, Bl. : Hook. Fil. Exot. t. 81.

A. Burchellii, Baker. Brazil.

A. buxifolium, Kze. *See* *sorbifolium*, var.

A. Cænopteris, Kze. Tropical America.

A. callæfolium, Bl. *See* *latifolium*, var.

A. cervinum, Sw. Tropical America. *Olfersia cervina.* Kze ; Hook. Fil. Exot. t. 43.

A. conforme, Sw. Tropical and sub-tropical regions.


A. crassinerve, Kze. *See* *latifolium*, var.

A. crinitum, L. West Indies and Central America. Hook. & Grev. Ic. Fil. t. 1.

A. crispatum, Wall. *See* *virens*, var.

A. cuspidatum, Willd. West Indies to Peru.

A. decoratum, Kze. West Indies, Guiana, Brazil and Peru.



Acrostichum, L.—cont.

- A. decurrens**, Desv. Philippines. *Gymnopteris decurrentis*, Hook. Gard. Ferns, t. 6.
- A. drynarioides**, Hook. Penang, Solomon Isles, New Guinea.
- A. Feei**, Bory. West Indies to Ecuador.
- A. flagelliferum**, Wall. Tropical Asia. Hook. & Grev. Ic. Fil. t. 23.
- A. gramineum**, Jenm. West Indies.
- A. Herminieri**, Bory. West Indies to Brazil.
- A. heteromorphum**, Kl. Colombia and Ecuador.
- A. hybridum**, Bory. Tropical Africa. Tropical America.
- A. latifolium**, Sw. Tropics. Hook. Fil. Exot. t. 42.
 var. **angustifolium**, Hort. Tropical America.
 var. **callæfolium**, (Bl.) Baker. Java.
 var. **conopodium**, Hort.
 var. **crassinerve**, (Kze) Baker. Brazil.
- A. Lingua**, Raddi. West Indies and Tropical America.
- A. longifolium**, Jacq. See latifolium.
- A. luridum**, Fée. Trinidad, Guiana.
- A. magnum**, Baker. British Guiana.
- A. melanopus**, Kze. Venezuela.
- A. muscosum**, Sw. West Indies, Tropical America.
- A. osmundaceum**, Hook. West Indies, Tropical America.
- A. peltatum**, Sw. West Indies, Tropical America.
- A. prieurianum**, Kl. See sorbifolium, var.
- A. quercifolium**, Retz. Tropical Asia. *Gymnopteris quercifolia*, Bernh.; Hook. Fil. Exot. t. 80.

Acrostichum, L.—*cont.*

A. repandum, Bl. Tropical Asia, Queensland and the Seychelles.

var. *quoyanum*, Baker. Moluccas.

A. rigidum, Wall. Malay Peninsula and Islands.

A. scandens, J. Sm. Tropical Asia, Queensland and Fiji.

A. Sieberi, Hook. & Grev. Ic. Fil. t. 237. Mascarene Islands.

A. simplex, Sw. West Indies, Tropical America.

A. sorbifolium, L. Tropics.

var. *buxifolium*, (Kze) Baker. Madagascar.

var. *prieurianum*, (Kl.) Baker. Guiana.

A. spicatum, L. Tropics of the Old World. *Hymenolepis spicata*, Presl; Hook. Fil. Exot. t. 78.

A. tomentosum, Bory. Bourbon.

A. undulatum, Willd. See *villosum*, var.

A. variabile, Hook. Tropical Asia.

A. villosum, Sw. Tropical America. Hook. & Grev. Ic. Fil. t. 95.

var. *undulatum*, Baker. West Indies and Mexico to Ecuador.

A. virens, Wall. Tropical Asia, Tropical Africa. Hook. & Grev. Ic. Fil. t. 221.

var. *crispatum*, Baker. India.

Actiniopteris, Link.

A. radiata, Link. Tropical Asia, Tropical Africa. Hook. Ic. Pl. t. 975.

var. *australis*, Baker. Mascarene Islands. Hook. Ic. Pl. t. 976.

Adiantopsis. See *Cheilanthes*.

Adiantum, L.

- A. *aethiopicum*, L. Tropical and south temp. zones. Hook. & Grev. Ic. Fil. t. 77.
 var. *assimile*, (Sw.) Baker. Australia.
 var. *aureum*, Hort.
- A. *affine*, Hook. See *diaphanum*.
- A. *amabile*, Moore. See *Moorei*.
- A. *aneitense*, Carruth. Aneiteum.
- A. *assimile*. Sw. See *aethiopicum*, var.
- A. *Balfourii*, Baker. Socotra. Hook. Ic. Pl. t. 1630.
- A. *Bausei*, Moore. Garden origin. Gard. Chron. 1879, xi. 456, figs. 69-70.
- A. *bellum*, Moore. Bermuda, Guiana. Gard. Chron. 1879, xi. 172, fig. 24.
- A. *bessonianum*, O'Brien. See *tenerum*, var.
- A. *Birkenheadii*, Moore. Garden origin.
- A. **Capillus-Veneris*, L. Tropical and temp. zones. Hook. Brit. Ferns, t. 41.
- A. *cardiochlæna*, Kze. See *polyphyllum*.
- A. *Catharinæ*, J. Sm. See *trapeziforme*, var.
- A. *caudatum*, L. Tropics. Hook. Exot. Fl. t. 104.
 var. *ciliatum*, Hort.
- A. *chrysophyllum*, Hort. See *aethiopicum*, var. *aureum*.
- A. *clæsianum*, L. Lind. & Rod. Brazil. Ill. Hort. 1894, 137, t. 9.
- A. *Collisii*, Moore. Garden origin.
- A. *colpodes*, Moore. Tropical America.
 var. *elegans*, Hort.

* For garden varieties, see p. 157.

Adiantum, L.—cont.

A. concinnum, H. B. & K. Tropical America.

var. *Flemingii*, *Hort.*

var. *gracile*, *Hort.*

var. *latum*, *Moore.*

A. crenatum, Willd. West Indies, Mexico.

A. cristatum, L. West Indies, Tropical America.

A. cultratum, J. Sm. See *trapeziforme*, var.

A. cuneatum, Langsd. & Fisch. Tropical America
Hook. & Grev. Ic. Fil. t. 30.

var. *aemulum*, (*Moore*) *Baker.*

var. *deflexum*, *Hort.*

var. *dissectum*, *Hort.*

var. *elegantissimum*, *Hort.*

var. *gracillimum*, (*Moore*) *Baker.*

var. *grandiceps*, *Hort.*

var. *grandis*, *Hort.*

var. *lambertianum*, *Hort.*

var. *Legrandi*, *Hort.*

var. *mundulum*, (*Moore*) *Baker.*

var. *Pacotti*, *Hort.*

var. *Reginæ*, *Hort.*

var. *schizophyllum*, *Hort.*

var. *strictum*, *Hort.*

var. *variegatum*, *Hort.*

A. curvatum, Kaulf. Brazil. Hook. Sp. Fil. ii. t. 84,
fig. C.

var. *brasiliense*, *Baker.* Brazil.

A. cyclosorum, Moore. Ecuador.

A. decorum, Moore. See *Wagneri.*

***Adiantum, L.*—cont.**

A. densus, Hort. Garden origin.

A. diaphanum, Bl. S. China, Malaya, Polynesia, N. Zealand and New S. Wales.

A. digitatum, Presl. Tropical America.

var. *palmatum, Baker.* *A. palmatum, Moore, Gard. Chron. 1877, vii. 40, fig. 5.*

var. *speciosum, (Hook.) Baker.*

A. dolabriforme, Hook. See *lunulatum, var.*

A. dolosum, Kze. Tropical America.

A. Edgeworthii, Hook. Northern India and China. Hook. Sp. Fil. ii. t. 81, fig. B.

A. elegans, Moore. Garden origin.

A. emarginatum, Bory. See *aethiopicum.*

A. excisum, Kze. Chili.

var. *Leyi, Hort.*

var. *multifidum, Hort.*

var. *nanum, Hort.*

A. farleyense, Moore. See *tenerum, var.*

A. fasciculatum, Hort. Garden origin.

A. Faulkneri, Hort. Garden origin.

A. Fergusoni, Moore. Ceylon.

A. formosum, R. Br. Temperate Australia and New Zealand. Hook. Sp. Fil. ii. t. 86, fig. B.

A. fragile, Sw. West Indies.

A. fulvum, Raoul. New Zealand, N. S. Wales, Fiji. Hook. Sp. Fil. ii. t. 85, fig. A.

A. Funckii, Hort. See *trapeziforme.*

A. glaucophyllum, Hook. Central America.

Adiantum, L.—cont.

- A. grandiceps**, Hort. *See cuneatum*, var.
- A. henslovianum**, Hook. f. Colombia, Peru, &c.
- A. heterophyllum**, Hort. Garden origin.
- A. hirtum**, Klotzsch. Tropical America. Hook. Sp. Fil. ii. t. 82, fig. A.
- A. hispidulum**, Sw. Tropics of the Old World.
var. **tenellum**, Moore. Australia.
- A. Hosei**, Baker. Borneo.
- A. Kaulfussii**, Kze. Tropical America.
- A. kerchovianum**, Hort. Garden origin.
- A. Lathomi**, Moore. *See tenerum*, var.
- A. Legrandi**, Hort. *See cuneatum*, var.
- A. lineatum**, Linden. Garden origin. Ill. Hort. 1895, 312, t. 44.
- A. lucidum**, Sw. West Indies, Tropical America. Hook. Sp. Fil. ii. t. 79, fig. C.
- A. luddemannianum**, Hort. Garden origin.
- A. lunulatum**, Burm. Tropics. Hook. & Grev. Ic. Fil. t. 104.
var. **dolabriiforme**, Baker. *A. dolabriiforme*, Hook. Ic. Pl. t. 191.
var. **fernandezianum**, Hort.
- A. macrophyllum**, Sw. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 132.
var. **albostriatum**, Hort.
var. **bipinnatum**, Jenm. Jamaica.
- A. magnificum**, Hort. *See Wagneri*.
- A. manicatum**, Hort. *See concinnum*.

Adiantum, L.—cont.

- A. matthewsianum**, Hook. *See* polyphyllum.
- A. Moorei**, Baker. Andes of Peru.
- A. moritzianum**, Link. *See* A. Capillus-Veneris.
- A. mundulum**, Moore. *See* cuneatum, var.
- A. neoguineense**, Moore. New Guinea.
- A. Novæ-Caledoniæ**, Keys. New Caledonia.
- A. obliquum**, Willd. West Indies. Tropical America. Hook. & Grev. Ic. Fil. t. 190.
- A. obtusum**, Desv. West Indies. Tropical America. Hook. & Grev. Ic. Fil. t. 188.
- A. Pacotti**, Hort. *See* excisum.
- A. patens**, Willd. Tropical America. Hook. Sp. Fil. ii. t. 87, fig. A.
- A. pellucidum**, Mart. & Gal. *See* æthiopicum.
- A. pedatum**, L. Temperate Asia. North America.
- A. peruvianum**, Klotzsch. Peru.
- A. plumosum**, Hort. Garden origin.
- A. polyphyllum**, Willd. Tropical America. Hook. Gard. Ferns, t. 12.
- A. prionophyllum**, H. B. & K. *See* tetraphyllum.
- A. pulverulentum**, L. West Indies, Tropical America.
- A. Reginæ**, Hort. *See* Capillus-Veneris.
- A. reniforme**, L. Madeira and Teneriffe. Hook. Fil. Exot. t. 8.
- var. **asarifolium**, Baker. Mauritius and Bourbon.
A. asarifolium, Willd. : Hook. Fil. Exot. t. 11.
- A. rhodophyllum**, Moore. Garden origin.
- A. rhomboideum**, Hort. *See* obtusum.

Adiantum, L.—cont.

- A. roseum**, Hort. Garden origin.
- A. rubellum**, Moore. Andes of Bolivia.
- A. scabrum**, Kze. See *aethiopicum*.
- A. scutum**, Hort. See *tenerum*, var.
- A. Seemanni**, Hook. Central America, Brazil. Gard. Chron. 1875, iii. 396–7, fig. 77–8.
- A. sessilifolium**, Hook. See *henslovianum*.
- A. setulosum**, J. Sm. See *diaphanum*.
- A. speciosum**, Hook. See *digitatum*, var.
- A. Tabarum**, Hort. See *excisum*.
- A. tenellum**, Moore. See *hispidulum*.
- A. tenerum**, Sw. West Indies, Tropical America.
 var. *alcicorne*, Hort.
 var. *bessonianum*, (O'Brien).
 var. *farleyense*, André. *A. farleyense*, Moore : Fl. Serres, tt. 1933–4.
 var. *gracile*, Hort.
 var. *Lathomi*, Hort.
 var. *Reginæ*, Hort.
 var. *scutum*, Hort.
 var. *scutum ramosum*, Hort.
 var. *Victoriæ*, (Moore) Baker.
- A. tetraphyllum**, Willd. West Indies, Tropical America.
 var. *acuminatum*, Hort.
 var. *gracile*, Hort.
- A. tinctum**, Moore. Andes of Peru.

Adiantum, L.—cont.

- A. trapeziforme, L.** West Indies, Tropical America.
 var. *Catharinæ, J. Sm.*
 var. *cultratum, Baker.*
 var. *Funckii, Hort.*
- A. veitchianum, Moore.** Andes of Peru.
- A. velutinum, Moore.** Andes of Colombia.
- A. venustum, G. Don.** Afghanistan and Himalaya.
 Hook. Sp. Fil. ii. t. 76, fig. B.
 var. *integrifolium, Baker.* Himalaya.
- A. Victoriae, Moore.** See *tenerum*, var.
- A. Wagneri, Mett.** Andes of Peru.
- A. Weigandii, Moore.** Garden origin.
- A. Williamsii, Moore.** Andes of Peru. Gard. Chron. 1878, x. 45, fig. 4.

Aglaomorpha. See *Polypodium*.

Aleuritopteris. See *Cheilanthes*.

Allosorus. See *Pellaea* and *Cryptogramme*.

Alsophila, R. Br.

- A. armata, Presl.** Tropical America.
- A. australis, R. Br.** Australia and Tasmania. Hook. Sp. Fil. i. t. 19, fig. A.
- A. contaminans, Wall.** India, Malaya. Fl. des Serres, tt. 2264–5.
- A. Cooperi, Hook.** Australia.
- A. excelsa, R. Br.** Norfolk Island.
- A. infesta, Kze.** Tropical America.

Alsophila, R. Br.—cont.

- A. leichardtiana**, *F. Muell.* Australia. *Flor. & Pom.* 1873, 229.
- A. MacArthuri**, Hook. *See* leichardtiana.
- A. Malzinei**, Hort. Mexico.
- A. nigra**, Jenm. (*an Mart.?*). Trinidad.
- A. procera**, Kaulf. Tropical America.
- A. pruinata**, Kaulf. West Indies, Tropical and Temperate S. America.
- A. robusta**, De Vriese. Java.
- A. Tænitis**, Hook. Brazil.
- A. tomentosa**, Hook. Java, Formosa.
- A. truncata**, Brack. Fiji, Samoa.

Amphicosmia. *See* Hemitelia.

Anapeltis. *See* Polypodium.

A. nitida, J. Sm. *See* P. lycopodioides.

Anchistea. *See* Woodwardia.

Anemia, Sw.

- A. adiantifolia**, Sw. West Indies, Tropical America
Hook & Grev. Ic. Fil. t. 16.
- A. collina**, Raddi. Brazil. Hook. Fil. Exot. t. 1.
- A. dregeana**, Kze. Natal. Hook. Ic. Pl. t. 236.
- A. frarinifolia**, Raddi. *See* Phyllitidis.
- A. fulva**, Sw. *See* tomentosa, var.
- A. hirsuta**, Sw. West Indies, Tropical America.
- A. longifolia**, Link. *See* Phyllitidis, var.

Anemia, Sw.—*cont.*

A. Phylltidis, Sw. West Indies, Tropical America.
Anemidictyon Phylltidis, J. Sm.; Hook. Gen. Fil. t. 103.

var. *fœtida*, Hort.

var. *longifolia*, Hort.

A. tomentosa, Sw. West Indies, Tropical America.
 Hook. Fil. Exot. t. 30.

var. *fulva*, Moore. Tropical America. *A. fulva*, Sw.; Hook. Fil. Exot. t. 26.

Anemidictyon. See Anemia.

Anetium. See Hemionitis.

Angiopteris, Hoffm.

A. evecta, Hoffm. Tropics of the Old World. Hook. Fil. Exot. t. 75.

var. *brongniartiana*, (De Vriese) Baker. Tahiti.

var. *durvilleana*, (De Vriese) Baker. Polynesia, Manilla.

var. *hypoleuca*, (De Vriese) Baker. Java.

var. *miqueliania*, (De Vriese) Baker. Java.

var. *pruinosa*, (Kze) Baker. Java.

var. *teysmanniana*, (De Vriese) Baker. Java. Gard. Chron. 1905, xxxvii. May 27, Suppl.

Anisogonium. See Asplenium.

Antigramme. See Scolopendrium.

Arthropteris. See Polypodium tenellum, Nephrodium albopunctatum, and Nephrolepis.

Aspidium, Sw.

A. acrostichoides, Sw. North America.

A. *aculeatum, Sw. Ubiquitous.

var. *angulare*, *Gren. & Godr.* Hook. Brit. Ferns, t. 12.

var. *lobatum*, *Hook.* Hook. Brit. Ferns, t. 10.

var. *moritzianum*, (*Kl.*) *Baker.* Colombia.

var. *setosum*, (*Wall.*) *Baker.* Himalaya.

var. *tsus-simense*, *Baker.* Japan.

A. amabile, Bl. India, Formosa, Japan.

A. angulare, *Willd.* See *aculeatum*, var.

A. angulatum, *J. Sm.* See *Polypodium angulatum*.

A. anomalum, Hook. & Arn. Ceylon. *Polypodium anomalum*, *Hook. & Arn.* in *Hook. Journ. Bot.* viii. (1856) t. 11.

A. argutum, *Kaulf.* See *Nephrodium rigidum*, var. *americanum*.

A. aristatum, Sw. Tropical Asia, Polynesia, N. S. Wales, Natal. *Polystichum aristatum*, *Presl*: *Hook. f. Fl. N. Zealand*, t. 78.

var. *coniifolium*, (*Wall.*) *Baker.*

var. *variegatum*, *Hort.*

A. atratum, *Wall.* See *Nephrodium hirtipes*.

A. auriculatum, Sw. India, Formosa, Malaya.

A. Boottii, *Tuckerm.* See *Nephrodium spinulosum*, var.

A. capense, Willd. Tropical America and S. temperate zone.

A. caryotideum, *Wall.* See *falcatum*, var.

A. coadunatum, *Hook. & Grev.* See *Nephrodium cicutarium*.

* For garden varieties, see p. 157.

Aspidium, Sw.—cont.

- A. concavum**, Hort. *See* *laserpitiifolium*.
- A. confluens**, Mett. *See* *Nephrodium*.
- A. coniifolium**, Wall. *See* *aristatum*, var.
- A. coriaceum**, Sw. *See* *capense*.
- A. erythrosorum**, Eaton. *See* *Nephrodium*.
- A. Fadyenii**, Mett. *See* *Fadyenia prolifera*.
- A. falcatum**, Sw. Japan to South Africa. Hook. Fil. Exot. t. 92.
 var. **caryotideum**, Baker. Northern India, South Africa. *A. caryotideum*, Wall.; Hook & Grev. Ic. Fil. t. 69.
- var. **caryotideum attenuatum**, Hort.
- var. **Fortunei**, (J. Sm.) Baker. Japan.
- var. **pendulum**, Hort.
- A. festinum**, Hance. *See* *laserpitiifolium*.
- A. fragrans**, Sw. *See* *Nephrodium*.
- A. frondosum**, Lowe. Madeira.
- A. irregulare**, Brackenr. *See* *Nephrodium latifolium*.
- A. laserpitiifolium**, Mett. China, Japan, Tonquin.
- A. lepidocaulon**, Hook. Japan. *Polystichum lepidocaulon*, T. Moore in Gard. Chron. 1875, iii. 203, fig. 36.
- A. lobatum**, Sw. *See* *aculeatum*, var.
- A. Lonchitis**, Sw. N. temp. zone. Hook. Brit. Ferns, t. 9.
- A. macrophyllum**, Sw. *See* *Nephrodium*.
- A. mohrioides**, Bory. Temperate S. America.
- A. munitum**, Kaulf. W. North America.
 var. **imbricans**, Hort.

Aspidium, Sw.—cont.

- A.** *nevadense*, Boiss. *See* *Nephrodium rigidum*.
- A.** *nevadense*, Eaton. *See* *Nephrodium nevadense*.
- A.** *Plumieri*, Presl. *See* *trifoliatum*, var.
- A.** *polymorphum*, Wall. *See* *Nephrodium*.
- A.** *proliferum*, R. Br. *See* *aculeatum*.
- A.** *pungens*, Kaulf. South Africa.
- A.** *quinquangulare*, Kze. *See* *Nephrodium decompositum*.
- A.** *repandum*, Willd. Tropical Asia.
- A.** *Standishii*, Hort. *See* *laserpitifolium*.
- A.** *subtriphyllum*, Hook. *See* *Nephrodium*.
- A.** *triangulum*, Sw. West Indies. Hook, Fil. Exot. t. 33.
var. *ilicifolium*, Féé.
var. *xiphoides*, Moore.
- A.** *trifoliatum*, Sw. Tropical America.
var. *Plumieri*, (Presl) Baker.
- A.** *tripteron*, Kze. Japan.
- A.** *tsus-simense*, Hook. *See* *aculeatum*, var.
- A.** *uliginosum*, Nyman. *See* *Nephrodium cristatum*, var.
- A.** *variolosum*, Wall. *See* *Nephrodium*.
- A.** *varium*, Sw. China, Japan.
- A.** *vestitum*, Sw. *See* *aculeatum*.

Asplenium, L.

- A.** *abscissum*, Willd. Tropical America.
- A.** *achilleæfolium*, Liebm. Mexico.

Asplenium, L.—cont.

- A. acutum**, Bory. *See* *Adiantum-nigrum*, var. *Onopteris*.
- A. Adiantum-nigrum**, L. North and South temp. zones. Hook. Brit. Ferns, t. 33.
var. *Onopteris*, *Christens.*
- A. alatum**, H. B. & K. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 136.
- A. angustifolium**, Mich. North America.
- A. anisophyllum**, Kze. Tropical and S. Africa, Tropical America.
- A. apicidens*, Moore. *See* *Vieillardii*, var.
- A. arborescens**, Mett. Mascarene Islands.
- A. arboreum**, Willd. West Indies, Tropical America.
- A. Arnottii**, Baker. Sandwich Islands.
- A. attenuatum**, R. Br. Australia. Hook. & Grev. Ic. Fil. t. 220.
var. *integrum*, Hort.
- A. aureum**, Link. *See* *Ceterach*, var.
- A. auritum**, Sw. India, Mascarene Is., Trop. America.
var. *macilentum*, Baker.
var. *sulcatum*, Baker.
- A. australasicum**, Hook. *See* *Nidus*, var.
- A. axillare**, Webb & Berth. *See* *umbrosum*.
- A. bantamense**, Baker. Tropical Asia.
- A. Baptistii**, Moore. *See* *Vieillardii*, var.
- A. Belangeri**, Kze. Malayan Islands. Hook. Fil. Exot. t. 41.
- A. bifidum**, Presl. *See* *lineatum*, var.

Asplenium, L.—cont.

- A. brachypterum, Kze.** Tropical and South Africa, Mascarene Islands. Hook. Fil. Exot. t. 44.
- A. Brownii, J. Sm.** See umbrosum.
- A. bulbiferum, Forst.** Tropical Asia, Mascarene Is., Australia. Hook. Ic. Pl. t. 423.
- var. *fabianum*, Baker.
- var. *Hillii*, Hort.
- var. *laxum*, Hook. f. New Zealand.
- var. *minus*, Baker. New Zealand.
- var. *pumilum*, Hort.
- var. *tripinnatum*, Hook. f. New Zealand.
- A. caryotæfolium, Hort.** See cuneatum.
- A. celtidifolium, Kze.** Tropical America.
- A. Ceterach, L.** Europe and temp. Asia. Hook. Brit. Ferns, t. 36.
- var. *aureum*, Baker. Madeira and Canary Is.
- A. cicutarium, Sw.** Tropical America.
- A. Colensoi, Hook. f.** New Zealand.
- A. compressum, Sw.** St. Helena. Hook. Fil. Exot. t. 76.
- A. contiguum, Kaulf.** Polynesia and Malaya.
- var. *fissum*, Hort.
- A. cordifolium, Mett.** Malaya. *Diplazium cordifolium*, Bl.; Hook. Ic. Pl. t. 936.
- A. crenulatum, Baker.** West Indies, Tropical America.
- A. cultrifolium, L.** West Indies, Tropical America.
- A. cuneatum, Lam.** Polynesia, Africa, Trop. America. Hook. & Grev. Ic. Fil. t. 189.
- A. Currori, Hook.** Tropical Africa.

Asplenium, L.—cont.

- A. decussatum, Sw.** Tropics of Old World.
- A. dimidiatum, Sw.** Tropical Africa, Trop. America.
- A. dimorphum, Kze.** Norfolk Island.
- A. ebeneum, Ait.** Temperate regions.
- A. elongatum, Sw.** See *tenerum*, var.
- A. erectum, Bory.** See *lunulatum*.
- A. esculentum, Presl.** Tropical Asia. *Anisogonium esculentum*, Presl; Bedd. Handb. Ferns Ind. 193.
var. *glabrum*, Hort.
var. *serampurens*e, Baker.
- A. fabianum, Hombr. & Jacquem.** See *bulbiferum*, var.
- A. falcatum, Lam.** Tropics and sub-tropics, Old World.
var. *erosum*, Hort.
- A. feejeense, Brack.** Fiji Islands.
- A. fernandezianum, Kze.** See *lunulatum*, var.
- A. *Filia-fœmina, Bernh.** Ubiquitous. Hook. Brit. Ferns, t. 35.
- A. flabellifolium, Cav.** Australia. *A. flabelliforme*, Hook. Exot. Fl. t. 208.
var. *Richardsoni*, Hort.
- A. flaccidum, Forst.** Australia, Polynesia, Natal.
- A. fœniculaceum, H. B. & K.** See *fragrans*, var.
- A. fontanum, Bernh.** North temp. zone. Hook. Brit. Ferns, t. 34.
- A. fragrans, Sw.** West Indies, Tropical America.
var. *fœniculaceum*, Baker. *A. fœniculaceum*, H. B. & K.; Hook. & Grev. Ic. Fil. t. 92.

* For garden varieties, see p. 165.

Asplenium, L.—cont.

- A. furcatum**, *Thunb.* Tropical and sub-tropical regions.
 var. *tripinnatum*, *Baker.* Natal.
- A. germanicum**, *Weiss.* Europe. Hook. Brit. Ferns, t. 27.
- A. gæringianum*, Mett. See *macrocarpum*, var.
- A. grandifolium**, *Sw.* West Indies, Tropical America.
- A. griffithianum**, *Hook.* Ic. Pl. t. 928. Himalaya.
- A. Halleri*, Willd. See *fontanum*.
- A. Hemionitis**, *Wall.* South Europe and North Africa. Bot. Mag. t. 4911.
- A. heterocarpum**, *Wall.* India, Malaya. Bedd. Ferns S. Ind. t. 131.
- A. heterophlebium**, Mett. Himalaya. *Anisogonium heterophlebium*, Bedd. Handb. Ferns Ind. t. 192.
- A. hians**, *Kze.* West Indies, Tropical America.
- A. incisum**, *Thunb.* China, Japan.
- A. lanceolatum**, *Huds.* Europe.
- A. lanceum**, *Thunb.* China, Japan, India. *A. subsinuatum*, Hook. & Grev. Ic. Fil. t. 27.
- A. lasiopteris*, Mett. See *Peterseni*.
- A. laxum*, R. Br. See *bulbiferum*, var.
- A. lineatum**, *Sw.* Mauritius.
 var. *bifidum*, *Baker.*
 var. *inæquale*, *Baker.*
- A. longissimum**, *Bl.* Malaya, Mauritius.
- A. lucidum*, Forst. See *obtusatum*, var.
- A. lunulatum**, *Sw.* Tropics and subtropics.
 var. *erectum*, *Baker.* Mascarene Islands, Tropical America.

Asplenium, L.—cont.**A. lunulatum, Sw.—cont.**

- var. *fernandezianum*, (*Kze*) *Baker*. Juan Fernandez.
- var. *pteropus*, *Baker*. Tropical America. *A. pteropus*, Kaulf. ; Hook. Sp. Fil. iii. t. 177.
- var. *tenellum*, (*Roxb.*) *Baker*. St. Helena.
- var. *tripinnatifidum*, *Hort*.

A. macilentum, *Kze*. See *auritum*, var.

A. macrocarpum, Bl. Tropical Asia.

- var. *gœringianum*, (*Mett.*) *Baker*. Japan.

A. marginatum, L. West Indies, Tropical America. Hook. Fil. Exot. t. 73.

A. marinum, L. Europe, Canaries and Azores. Hook. Brit. Ferns, t. 31.

- var. *imbricatum*, *Hort*.

- var. *ramosum*, *Hort*.

A. maximum, G. Don. Tropical Asia, Australia

A. Mayii, Hort. Garden origin. Gard. Chron. 1898, xxiii. 332, fig. 140.

A. Michauxii, Spreng. See *Filix-fœmina*.

A. monanthemum, L. Temperate regions.

A. Nidus, L. Tropics, Old World. Bot. Mag. t. 3101.

- var. *australasicum*, *Baker*. Australia. *A. australasicum*, Hook. Fil. Exot. t. 88.

- var. *madagascariense*, *Hort*.

- var. *multilobatum*, *Hort*.

A. niponicum, Mett. Japan.

- var. *cristatum*, *Hort*.

A. nitidum, Sw. India and Malaya.

A. nobile, Hort. See *viviparum*.

A. nodulosum, Hort. See *lineatum*.

Asplenium, L.—cont.

- A. obliquum**, Forst. *See* obtusatum, var.
- A. obtusatum**, Forsi. South temperate regions. Hook.
Fil. Exot. t. 46.
- var. *lucidum*, Baker.
- var. *Lyallii*, Hook. New Zealand.
- var. *obliquum*, Hook. f. New Zealand.
- A. obtusilobum**, Hook. Polynesia.
- A. ornatum**, Col. New Zealand.
- A. paleaceum**, R. Br. Tropical Australia. Hook. Sp.
Fil. iii. t. 199.
- A. Peterseni**, Kze. India, Java.
- A. planicaule**, Wall. India, China.
- A. plantagineum**, L. West Indies, Tropical America.
- A. platyneuron**, Oakes. *See* ebeneum.
- A. polyodon**, Forst. *See* falcatum.
- A. polypodioides**, Mett. Tropical Asia. *Diplazium polypodioides*, Bl. ; Bedd. Handb. Ferns Ind. 185.
- A. præmorsum**, Sw. *See* furcatum.
- A. procerum**, Wall. Northern India.
- A. proliferum**, Lam. *See* decussatum.
- A. prolongatum**, Hook. *See* rutæfolium.
- A. pteridoides**, Baker. Lord Howe's Island.
- A. pteropus**, Kaulf. *See* lunulatum, var.
- A. pubescens**, Mett. *See* esculentum.
- A. pumilum**, Sw. West Indies, Tropical America.
- A. pusillum**, Bl. Java.
- A. radicans**, Schkuhr. West Indies, Tropical America.

Asplenium, L.—cont.

- A. reclinatum**, Houlst. *See lunulatum*, var. *tenellum*.
- A. refractum**, Moore. *See fontanum*.
- A. resectum**, Sm. Japan, Tropical Asia and Africa. Hook. & Grev. Ic. Fil. t. 114.
- A. rhizophorum**, L. West Indies, Tropical America. Hook. Sp. Fil. iii. t. 187.
- A. rigidum**, Sw. *See auritum*, var. *sulcatum*.
- A. rutæfolium**, Kze. Tropical and sub-tropical Asia and Africa.
var. *prolongatum*, (Hook.) Baker.
- A. Ruta-muraria**, L. North temp. zone. Hook. Brit. Ferns, t. 28.
- A. scandens**, J. Sm. New Guinea, Philippines, Fiji.
- A. semihastatum**, Kze. Cuba.
- A. septentrionale**, Hoffm. North temp. zone. Hook. Brit. Ferns, t. 26.
- A. Serra**, Langsd. & Fisch. West Indies, Tropical America and Africa.
var. *natalensis*, Baker. South Africa.
- A. Sheperdi**, Spreng. West Indies, Tropical America.
- A. sinuatum**, Beauv. Tropical Africa. Hook. Fil. Exot. t. 61.
- A. speciosum**, Mett. Malay Peninsula and Archipelago.
- A. striatum**, L. *See Sheperdi*.
- A. subsinuatum**, Hook. & Grev. *See lanceum*.
- A. sylvaticum**, Presl. Tropics, Old World.
- A. tenellum**, Roxb. *See lunulatum*, var.
- A. tenerum**, Forst. Ceylon, Malaya, Polynesia.
var. *elongatum*, (Sw.) Baker. Malaya.

Asplenium, L.—cont.


- A. *thelypteroides*, *Mich.* Temp. East Asia, North America.
- A. *tomentosum*, *Hook.* N. India, Malay Peninsula and Islands.
- A. *Trichomanes*, *L.* Temp. regions. *Hook. Brit. Ferns*, t. 29.
- A. *umbrosum*, *J. Sm.* Tropical Asia, Africa and Australia, Canary Islands.
var. *procerum*, Baker. *See procerum.*
- A. *unilaterale*, Lam. *See resectum.*
- A. *veitchianum*, Moore. *See Belangeri.*
- A. *Vieillardii*, Mett. New Caledonia.
var. *apicidens*, (*Moore*) Baker. South Sea Islands.
var. *Baptistii*, (*Moore*) Baker. South Sea Islands.
var. *Drueryi*, Hort.
- A. *virescens*, Mett. Japan.
- A. *viride*, *Huds.* North temp. zone. *Hook. Brit. Ferns*, t. 30.
- A. *viviparum*, *Presl.* Mascarene Islands. *Hook. Fil. Exot.* t. 64.
var. *nobile*, Hort.
- A. *vulcanicum*, Bl. Malay Islands.
- A. *zeylanicum*, *Hook.* Ceylon. *Sec. Cent. Ferns*, t. 16.

Athyrium. *See Asplenium.*

Balantium. *See Dicksonia.*

Blechnum, L.

- B. *Atherstonei*, Pappe. *See Lomaria punctulata*, var. *blechnoides*.



Blechnum, L.—cont.

- B. Banksii*, Mett. *See Lomaria.*
- B. boreale*, Sw. *See Lomaria Spicant.*
- B. brasiliense*, Desv. Tropical America.
var. *corcovadense*, (Raddi) Baker. Brazil.
var. *crispum*, Hort.
- B. cartilagineum*, Sw. Temperate Australia.
- B. corcovadense*, Raddi. *See brasiliense*, var.
- B. Gilliesii*, Mett. *See Lomaria procera.*
- B. glandulosum*, Link. *See occidentale*, var. *distans*.
- B. gracile*, Kaulf. *See longifolium*, var.
- B. hastatum*, Kaulf. Temperate South America. *B. trilobum*, Presl; Hook. & Grev. Ic. Fil. t. 192.
- B. Lanceola*, Sw. Tropical America. Bot. Mag. t. 3240.
- B. latifolium*, Hort. *See longifolium.*
- B. lomarioides*, Gaud. *See orientale.*
- B. longifolium*, H. B. & K. Tropical America.
var. *fraxineum*, Baker.
var. *gracile*, Baker.
- B. occidentale*, L. West Indies, Trop. America, Chili.
var. *distans*, Baker. S. Brazil, Uruguay.
var. *cristatum*, Hort.
- B. orientale*, L. S. China to Polynesia and Australia.
Hook. Fil. Exot. t. 77.
- B. polypodioides*, Raddi. *See unilaterale.*
- B. punctulatum*, Sw. *See Lomaria.*
- B. rugosum*, Moore. Origin unknown.
- B. triangulare*, Link. *See unilaterale.*

Blechnum, L.—*cont.*

B. trilobum, Presl. *See hastatum.*

B. unilaterale, Willd. West Indies, Tropical America.

Bolbitis. *See Acrostichum.*

Botrychium, Sw.

B. ternatum, Sw. North and south temp. zones.

var. *dissectum*, Baker. North America, New Zealand.

B. virginianum, Sw. North temp. zone, Trop. America. Hook. Gard. Ferns, t. 29.

Brainea, Hook.

B. insignis, Hook. Fil. Exot. t. 38. Hong Kong and Khasia.

Cænopteris. *See Asplenium.*

Callipteris. *See Asplenium.*

Campyloneuron. *See Polypodium.*

Campteria. *See Pteris.*

Camptosorus. *See Scolopendrium.*

Ceratopteris, Brongn.

C. thalictroides, Brongn. Tropics. *Parkeria pteridoides*, Hook & Grev. Ic. Fil. t. 97; Hook. Exot. Fl. iii. t. 231.

Ceropteris. *See Gymnogramme.*

Ceterach. *See Asplenium.*

Cheilanthes, Sw.

- C. anceps**, Blandf. *See* farinosa, var.
- C. argentea**, Hook. Fil. Exot. t. 95. North Asia, Malay Peninsula.
- C. dicksonioides**, Endl. *See* Hypolepis tenuifolia.
- C. elegans**, Desv. *See* myriophylla, var.
- C. ellisiana**, Hort. *See* hirta, var.
- C. farinosa**, Kaulf. Tropics. Bot. Mag. t. 4765.
var. **anceps**, Blandf. Himalaya.
var. **punctata**, Hort.
- C. fragrans**, Sw. Europe, Orient, N. Africa, Canary Is.
- C. hirta**, Sw. Tropical and South Africa.
var. **ellisiana**, Hort.
- C. mexicana**, Davenp. Mexico.
- C. micromera**, Link. *See* microphylla, var.
- C. microphylla**, Sw. West Indies, Tropical America.
var. **micromera**, Baker. Central America.
- C. myriophylla**, Desv. Tropical America.
var. **elegans**, Baker.
- C. profusa**, Kze. *See* Pellaea involuta.
- C. radiata**, R. Br. West Indies, Tropical America.
- C. tomentosa**, Link. United States and Mexico.
- C. undulata**, Hope & Wright. China.
- C. vestita**, Sw. United States.

Chrysodium. *See* Acrostichum.

Cibotium. *See* Dicksonia.

Cincinalis. *See* *Notholæna.*

Cionidium. *See* *Deparia.*

Coniogramme. *See* *Gymnogramme.*

Conpteris. *See* *Lonchitis.*

C. brazziana, Linden. *See* *Lonchitis pubescens*, var.

Craspedaria. *See* *Polypodium.*

Cryptogramme, R. Br.

C. crispa, *R. Br.* North temp. zone. Hook. Brit. Ferns, t. 39.

var. *acrostichoides*, *Baker.* North America. *C. acrostichoides*, *R. Br.*; Hook & Grev. Ic. Fil. t. 29.

Cyathea, J. E. Smith.

C. arborea, *J. E. Sm.* West Indies, Tropical America.

C. beyrichiana, Presl. *See* *Hemitelia setosa.*

C. dealbata, *Sw.* New Zealand.

var. *tricolor*, (*Col.*) *Baker.*

C. Dregei, *Kze.* Tropical and South Africa.

var. *Burkei*, *Hook.* Tropical and South Africa.

C. excelsa, *Sw.* Bourbon and Mauritius.

C. gracilis, *Griseb.* Jamaica.


C. Hookeri, *Thw.* Ceylon.

C. imrayana, *Hook.* *See* *Tussacii.*

C. insignis, *D. C. Eaton.* West Indies.

C. medullaris, *Sw.* New Zealand. Hook. Gard. Ferns, t. 35.

*Cyathea, J. E. Smith—cont.**C. mexicana, Schlecht.* Mexico.*C. princeps, J. Sm.* See *insignis*.*C. Serra, Willd.* West Indies, Tropical America.*C. Smithii, Hook. f.* See *Hemitelia Smithii*.*C. spinulosa, Wall.* India, Japan.*C. tricolor, Col.* See *dealbata*.*C. Tussacii, Desv.* West Indies.*Cyclodium.* See *Aspidium*.*Cyclopeltis.* See *Aspidium*.*Cyrtogonium.* See *Acrostichum*.*Cyrtomium.* See *Aspidium*.*Cyrtophlebium.* See *Polypodium*.*Cystopteris, Bernh.**C. alpina, Desv.* Europe, Asia Minor. Hook. Brit. Ferns, t. 24.*C. bulbifera, Bernh.* North America.*C. fragilis, Bernh.* Cosmopolitan. Hook. Brit. Ferns, t. 23.var. *angustata, Link.* Europe.var. *dickieana, Moore.* Scotland.*C. montana, Link.* Europe, North America. Hook. Brit. Ferns, t. 25.*C. tenuis, Schott.* See *fragilis*.*Darea.* See *Asplenium*.



Davallia, J. E. Smith.

D. aculeata, *Sw.* West Indies.

var. *inermis*, *Hort.*

D. affinis, *Hook.* Tropical and sub-tropical Asia, Polynesia.

D. alpina, *Bl.* Malaya, Polynesia.

D. angustata, *Wall.* Malay Peninsula and Islands. *Hook. & Grev. Ic. Fil. t. 231.*

D. assamica, *Baker.* Bhotan, Manipur.

D. bullata, *Wall.* Japan, Tropical Asia.

var. *cristata*, *Hort.*

var. *cypnochlamys*, *Baker & Clarke.* Northern India.

var. *decora*, (*Moore*) *Baker.* Java.

var. *Mariesii*, (*Moore*) *Baker.* Japan.

D. canariensis, *J. E. Sm.* Western Europe and the Canaries.

D. chærophylla, *Wall.* India, Malaya.

D. ciliata, *Hook.* Philippines.

var. *angustata*, *Hort.*

D. decora, *Moore.* See *bullata*, var.

D. decurrens, *Hook.* Philippines.

D. dissecta, *J. Sm.* Java.

var. *elegans*, *Hort.*

D. divaricata, *Bl.* India, Malaya.

D. elata, *Sw.* See *elegans*, var.

D. elegans, *Sw.* Tropical Asia, Africa and Australia.

var. *elata*, *Baker.*

var. *Kunzei*, *Hort.* Malya.

var. *polydactyla*, *Hort.*

Davallia, J. E. Smith—cont.

- D. epiphylla**, Bl. Malay Peninsula, Java.
- D. fejeensis**, Hook. Fiji Islands.
- var. **effusa**, Hort.
- var. **gracillima**, Hort.
- var. **major**, Hort.
- var. **plumosa**, Hort.
- var. **robusta**, Hort.
- D. fijiensis**, Hook. & Baker. *See fejeensis.*
- D. fœniculacea**, Hook. Fiji.
- D. glabella**, Hort. Garden origin.
- D. griffithiana**, Hook. China, India, Malaya.
- D. hemiptera**, Bory. *See repens.*
- D. heterophylla**, J. E. Sm. Malay Peninsula, Polynesia.
Hook. Fil. Exot. t. 27.
- D. hirta**, Kaulf. India, Malaya, Polynesia.
var. **cristata**, Hort.
- D. illustris**, Hort. Garden origin.
- D. immersa**, Wall. India, Malaya. Hook. Fil. Exot.
t. 79.
- D. khasiana**, Hook. *See strigosa.*
- D. lawsoniana**, Hort. Garden origin.
- D. Lindleyi**, Hook. *See solida*, var. *caudata.*
- D. Lorranei**, Hance. Malay Peninsula.
- D. lonchitidea**, Wall. *See platyphylla.*
- D. lucida**, Hort. *See solida.*
- D. madagascariensis**, Baker. Madagascar.
- D. majuscula**, Lowe. Ceylon.

Davallia, J. E. Smith—cont.

- D. marginalis, Baker.** China, Japan, India.
- D. Mariesii, Moore.** *See* bullata, var.
- D. mauritiana, Hook.** Mauritius.
- D. mooreana, Masters.** *See* pallida.
- D. nitidula, Kze.** Tropical and South Africa.
- D. Novæ-Zealandiæ, Col.** New Zealand.
- D. ornata, Wall.** *See* solida, var. latifolia.
- D. pallida, Mett.** Malaya and Polynesia.
var. *intermedia*, Hort.
- D. parvula, Wall.** Malay Peninsula and Archipelago.
- D. pedata, J. E. Sm.** Tropical and subtropical Asia, Australia, Mascarene Islands. Hook. Gard. Ferns, t. 7.
- D. pentaphylla, Bl.** Java, Polynesia. Hook. Fil. Exot. t. 37.
- D. platyphylla, D. Don.** India. *D. lonchitidea*, Wall.; Hook. Fil. Exot. t. 19.
- D. polyantha, Hook.** *See* divaricata.
- D. polypodioides, D. Don.** *See* Speluncæ.
- D. pulchella, Hook.** Philippines, Fiji, Samoa.
- D. pulchra, D. Don.** China, India.
- D. pycnocarpa, Brackenr.** *See* pentaphylla.
- D. repens, Desv.** India, Malaya, Polynesia.
- D. retusa, Cav.** *See* Lindsaya retusa.
- D. rhomboidea, Wall.** *See* strigosa, var.
- D. scabra, D. Don.** *See* marginalis.

Davallia, J. E. Smith—cont.

D. solida, Sw. Malaya, Polynesia, Queensland. Hook. Fil. Exot. t. 57.

var. **caudata**, Hook. Singapore.

var. **latifolia**, Hook. Penang.

var. *Lindleyi*, Hook. See var. caudata.

var. **lucida**, Hort.

D. Speluncæ, Baker. Tropics, ubiquitous.

D. stenocarpa, Hort. Par. See *madagascariensis*.

D. strigosa, Sw. Tropical and subtropical Asia, Polynesia.

var. **rhomboidea**, Hook. Northern India.

D. tenuifolia, Sw. Tropics of the Old World.

var. **stricta**, Hort.

var. **Veitchii**, Hort.

D. trichosticha, Hook. See *Speluncæ*.

D. triphylla, Hook. Singapore.

D. Tyermannii, Baker. China. Hook. Ic. Pl. t. 1620.

D. villosa, Wall. See *marginalis*.

Dennstædtia. See *Dicksonia*.

Deparia, Hook. & Grev.

D. Moorei, Hook. New Caledonia. Hook. Fil. Exot. t. 28.

Diacalpe, Bl.

D. aspidioides, Bl. Tropical Asia.

Dicksonia, L'Hérit.

D. adiantoides, H. B. & K. West Indies, Tropical America.

Dicksonia, L'Hérit.—cont.

- D. antarctica**, *Labill.* Australia.
- D. apiifolia**, *Hook.* Andes of Ecuador and Peru.
- D. arborescens**, *L'Hérit.* St. Helena.
- D. Barometz**, *Link.* Sub-tropical Asia.
- D. Chamissoi**, *Hook. & Baker.* Sandwich Islands.
- D. cicutaria**, *Sw.* West Indies, Tropical America.
var. **tenera**, *Hort.*
- D. Culcita**, *L'Hérit.* Madeira, Azores and Spain.
- D. davallioides**, *R. Br.* Australia.
- D. Deplanchei**, *Vieill.* New Caledonia.
- D. fibrosa**, *Col.* New Zealand.
- D. glauca**, *J. E. Sm.* Sandwich Islands.
- D. gracilis**, *Col.* New Zealand.
- D. Herberti**, *Hill.* Queensland ?.
- D. karsteniana*, Klotzsch. See *sellowiana*, var.
- D. Lathami**, *Moore.* Garden origin. Gard. Chron. 1885,
xxiv. 584, 689, fig. 154.
- D. Menziesii**, *Hook. & Baker.* Sandwich Islands.
- D. obtusifolia*, Willd. See *adiantoides*.
- D. princeps*, Hort. See *Cyathea insignis*.
- D. punctiloba*, Hook. See *punctilobula*.
- D. punctilobula*, *A. Gray.* North America.
- D. regalis**, *Baker.* Mexico.
- D. rubiginosa**, *Kaulf.* West Indies, Tropical America.
- D. Schiedei**, *Baker.* Central America.

Dicksonia, L'Hérit.—cont.

D. sellowiana, Hook. South Brazil.

var. *karsteniana, Baker.* Guatemala and Andes of Colombia.

D. spectabilis, Hort. See *Wendlandi*.

D. squarrosa, Sw. New Zealand.

D. tenera, Hort. See *apiifolia*.

D. Wendlandi, Baker. Guatemala.

var. *Verschaffeltii, Hort.*

D. Youngiae, C. Moore. New South Wales.

Dictyogramme. See *Gymnogramme*.

Dictyopteris. See *Polypodium*.

Didymochlæna, Desv.

D. lunulata, Desv. Tropics, ubiquitous. Hook. Gard. Ferns, t. 17.

D. sinuosa, Desv. See *lunulata*.

D. truncatula, J. Sm. See *lunulata*.

Didymoglossum. See *Hymenophyllum*.

Diplazium. See *Asplenium*.

Dipteris. See *Polypodium*.

Disphenia. See *Cyathea*.

Doodia, R. Br.

D. amœna, Hort. Garden origin.

D. aspera, R. Br. Temperate Australia.

var. *multifida, Hort.*

Doodia, R. Br.—cont.

- D. blechnoides**, Cunn. Australia.
D. caudata, R. Br. Australia and New Zealand.
D. kunthiana, Gaud. Sandwich Islands.
D. lunulata, R. Br. *See media.*
D. media, R. Br. Polynesia, Australia, New Zealand.

Doryopteris. *See* Pteris.**Drymoglossum, Presl.**

- D. carnosum**, Hook. Eastern Himalaya.
var. **minor**, Hook. China and Japan.
D. piloselloides, Presl. India, Malaya. Hook. Gard. Ferns, t. 46.
D. spathulatum, Presl. *See* piloselloides.
D. subcordatum, Fée. *See* carnosum, var. minor

Drynaria. *See* Polypodium.

- D. diversifolia**, J. Sm. *See* P. rigidulum.
D. morbillosa, J. Sm. *See* P. Heracleum.

Dryopteris. *See* Nephrodium.

- D. acrostichoides**, O. Kze. *See* Aspidium.

Egenolfia. *See* Acrostichum.**Elaphoglossum. *See* Acrostichum.****Eupodium. *See* Marattia.****Fadyenia, Hook.**

- F. prolifera**, Hook. West Indies.

Feea. *See Trichomanes.*

Gleichenia, J. E. Smith.

G. circinata, Sw. Australia, New Zealand, Malaya.

var. *backhousiana*, *Hort.*

var. *semivestita*, (*Labill.*) *Baker.* New Caledonia.

G. dicarpa, R. Br. Australia, New Zealand. Hook.

Fil. Exot. t. 40.

var. *alpina*, *Hook.* Tasmania, New Zealand, Malaya.

var. *glaucia*, *Moore.*

var. *longipinnata*, *Hort.*

G. flabellata, R. Br. Australia, New Zealand, New

Caledonia. Hook. Fil. Exot. t. 71.

G. longipinnata, *Hort.* *See dicarpa, var.*

G. rupestris, R. Br. Australia.

var. *glaucescens*, *Hort.*

Goniophlebium. *See Polypodium.*

Goniopteris. *See Polypodium.*

Gymnogramme, Desv.

G. ampla, Benth. Australia.

G. calomelanos, Kaulf. West Indies, Tropical America.
Hook. Gard. Ferns, t. 50.

var. *Alstoni*, *Hort.*

var. *argyrophylla*, *Hort.*


var. *chrysophylla*, *Baker.*

var. *grandiceps*, *Hort.*

var. *grandiceps superba*, *Hort.*

var. *intermedia*, (*Fée*).

var. *Martensii*, (*Link*) *Baker.*



Gymnogramme, Desv.—cont.

- G. caudiformis, Hook.** Tropical Asia, Polynesia. *Grammitis caudiformis*, Hook. Bot. Mag. t. 5328.
- G. chrysophylla, Kaulf.** See calomelanos.
- G. decurrens, Hook.** See elliptica.
- G. dobroydense, Hort.** Garden origin.
var. **cristata, Hort.**
- G. elliptica, Baker.** Tropical Asia, Queensland. *Grammitis decurrens*, Hook. & Grev. Ic. Fil. t. 6.
- G. gracilis, Heward.** Jamaica.
- G. hamiltoniana, Hook.** Eastern Himalaya.
- G. Heyderi, Lauche.** Dominica.
var. **argentea, Hort.**
- G. japonica, Hook.** Japan, Formosa.
- G. laucheana, K. Koch.** See calomelanos.
- G. leptophylla, Desv.** Cosmopolitan. Hook. Brit. Ferns, t. 1.
- G. L'Herminieri, Link.** See calomelanos.
- G. Mathewsii, Hook.** Peru.
- G. Muelleri, Hook.** Australia.
- G. multiceps, Hort.** See calomelanos.
- G. ochracea, Presl.** See tartarea, var.
- G. Parsonsii, Moore.** See calomelanos.
- G. Pearcei, Moore.** Peru.
var. **robusta, Hort.**

Gymnogramme, Desv.—cont.

G. peruviana, Desv. *See calomelanos.*

G. pulchella, Linden. Venezuela. Hook. Fil. Exot. t. 74.

var. *wettenhalliana*, Hort.

var. *Mayi*, Hort.

G. rufa, Desv. West Indies, Tropical America.

G. schizophylla, Baker. Hook. Ic. Pl. t. 1682.

var. *elegantissima*, Hort.

var. *gloriosa*, Rodigas. Ill. Hort. 1884, t. 522.

G. sulphurea, Desv. West Indies.

var. *grandiceps*, Hort.

G. tartarea, Desv. Tropical America.

var. *ochracea*, Baker. South Brazil.

G. tomentosa, Desv. South America. Hook. Fil. Exot. t. 13.

G. triangularis, Kaulf. From Vancouver Island to Ecuador. Hook. Fil. Exot. t. 10.

G. Veitchii, Hort. Garden hybrid.

Gymnopteris. See Acrostichum.**Haplopteris. See Vittaria.****Helminthostachys, Kaulf.**

H. dulcis, Kaulf. *See zeylanica.*

H. zeylanica, Hook. India, Malaya, Queensland. Hook. Gard. Ferns, t. 28.

Hemidictyum. See Asplenium.

Hemionitis, L.

H. cordata, Roxb. India, Malaya. Hook. & Grev. Ic. Fil. t. 64.

H. palmata, L. West Indies, Tropical America.

Hemitelia, R. Br.

H. capensis, R. Br. South Africa, Brazil.

H. setosa, Mett. Brazil.

H. Smithii, Hook. New Zealand.

H. speciosa, Hook. Colombia. Hook. Fil. Exot. t. 66.
var. *Klotzschii*, Hort.

Humata. See Davallia.

Hymenodium. See Acrostichum.

Hymenolepis. See Acrostichum.

Hymenophyllum, J. E. Sm.

H. asplenoides, Sw. West Indies, Tropical America.

H. axillare, Sw. West Indies, Venezuela.

H. capillaceum, Roxb. St. Helena.

H. Catharinæ, Hook. West Indies.

H. caudiculatum, Mart. Tropical America, Chili.

H. chiloense, Hook. South Chili and Island of Chiloe.

H. ciliatum, Sw. Tropical Africa and America. Hook. & Grev. Ic. Fil. t. 35.

H. crispatum, Wall. See *javanicum*, var.

H. cruentum, Cav. Chili. Hook. Sp. Fil. i. t. 31, fig. A.

Hymenophyllum, J. E. Sm.—cont.

- ✓ *H. demissum*, Sw. New Zealand, Polynesia, Malaya.
var. *nitens*, Hort.
- H. dilatatum*, Sw. New Zealand, Polynesia, Java.
Hook. & Grev. Ic. Fil. t. 60.
var. *fosterianum*, Hort.
- H. flabellatum*, Labill. Australia and New Zealand.
- H. flexuosum*, Cunn. See *javanicum*, var.
- H. fosterianum*, Hort. See *dilatatum*, var.
- H. fucoides*, Sw. West Indies, Tropical America. Hook.
Ic. Pl. t. 963.
- H. gracile*, Bory. South Africa, Mascarene Islands.
Hook. & Grev. Ic. Fil. t. 198.
- H. hirsutum*, Sw. West Indies, Tropical and Temperate
South America, Mascarene Islands. Hook. & Grev.
Ic. Fil. t. 84.
- H. hirtellum*, Sw. West Indies, Mexico.
- H. javanicum*, Spreng. Tropical Asia, Australia and
New Zealand.
var. *crispatum*, (Wall.) Baker. India.
var. *flexuosum*, (A. Cunn.) Baker. Australia. *H. flexuosum*, A. Cunn.; Hook. Ic. Pl. t. 962.
var. *montanum*, Hort.
var. *tamariscifolium*, Hort.
- H. lineare*, Sw. West Indies, Tropical America.
var. *elegantissima*, Hort.
- H. multifidum*, Sw. New Zealand, Polynesia.
- H. pectinatum*, Cav. Chili and Chiloe.
var. *superbum*, Hort.
- H. polyanthos*, Sw. Tropics, ubiquitous.
var. *villosum*, (Col.) Baker. New Zealand.

Hymenophyllum, J. E. Sm.—cont.

H. pulcherrimum, Col. New Zealand.

H. rarum, R. Br. S. Africa, Mascarene Islands, New Zealand.

H. scabrum, A. Rich. New Zealand.

H. sericeum, Sw. West Indies, Tropical America.

H. subtilissimum, Kze. New Zealand.

H. tamariscifolium, Hort. See *javanicum*, var.

H. tunbridgense, Sw. Ubiquitous. Hook. Brit. Ferns, t. 43.

var. *Wilsoni, Hook.* Britain, Norway. *H. Wilsoni, Hook.* Brit. Ferns, t. 44.

H. villosum, Col. See *polyanthos*, var.

Hypoderris, R. Br.

H. Brownii, J. Sm. Trinidad. Hook. Gard. Ferns, t. 24.

Hypolepis, Bernh.

H. bergiana, Hook. Tropical and South Africa.

H. millefolia, Hook. New Zealand.

H. purdieana, Hook. Colombia.

H. radiata, Hook. See *Cheilanthes*.


H. repens, Presl. West Indies, Tropical America.

H. tenuifolia, Bernh. Malaya, Polynesia, Australasia.

Isoloma. See *Lindsaya*.

Kaulfussia, Bl.

K. æsculifolia, Bl. Assam, Malaya. Hook. & Grev. Ic. Fil. t. 229.



Lastrea. *See Nephrodium.*

L. aristata, Hort. *See Aspidium aristatum.*

L. atrovirens, J. Sm. *See Nephrodium decompositum.*

L. opaca, Hook. *See Aspidium varium.*

Lepicystis. *See Polypodium.*

Leptochilus. *See Acrostichum.*

Leptogramme. *See Gymnogramme.*

Leptopteris. *See Todea.*

Leucostegia. *See Davallia.*

Lindsaya, Dry.

L. guianensis, Dry. West Indies, Tropical America.

L. linearis, Sw. Australia, New Zealand.

L. Lowii, Hort. *See Acrostichum sorbifolium.*

L. retusa, Mett. Malaya.

L. trapeziformis, Dry. Tropical America, Malaya.

var. *falcata, Baker.* West Indies.

Litobrochia. *See Pteris.*

Llavea, Lag.

L. cordifolia, Lag. Mexico. Bot. Mag. t. 5159.

Lomaria, Willd.

L. alpina, Spreng. South temperate zone. Hook. Fil. Exot. t. 32.

Lomaria, Willd—cont.

L. attenuata, Willd. Tropical and South temperate regions.

var. **gigantea**, (Kaulf.) Baker. South Africa. *L. gigantea*, Kaulf. ; Ill. Hort. 1876, t. 237.

var. **heterophylla**, Hort.

L. Banksii, Hook. f. New Zealand.

L. boryana, Willd. Tropical and South temperate regions.

L. capensis, Willd. See procera.

L. ciliata, Moore. New Caledonia. Ill. Hort. 1876, t. 252.

L. discolor, Willd. Australia and New Zealand.

var. **bipinnatifida**, Hort.

var. **nuda**, (R. Br.) Baker. Australia.

L. fluviatilis, Spreng. Australia and New Zealand. Hook. f. Fl. Tasm. t. 16.

L. gibba, Labill. Sert. Austr. Caled. tt. 4-5. New Caledonia, &c. *L. neocalledonica*, Linden & Fourn. Ill. Hort. 1876, t. 251.

var. **blechnoides**, Hort.

var. **tincta**, Hort.

L. gigantea, Kaulf. See attenuata.

L. Gilliesii, Hook. & Grev. See procera.

L. heterophylla, Hort. See attenuata.

L. lanceolata, Spreng. Australia, New Zealand, Polynesia. Hook. Ic. Pl. t. 429.

L. L'Herminieri, Bory. West Indies, Tropical America, Chili. Hook. Gard. Ferns, t. 40.

L. magellanica, Desv. See boryana.

Lomaria, Willd—cont.

L. membranacea, Col. New Zealand. Hook. Sp. Fil. iii. t. 145.

L. minor, R. Br. *See* procera.

L. neocaledonica, Linden & Fourr. *See* gibba.

L. nipponica, Kunze. Japan.

L. Patersonii, Spreng. Australia. Hook. Fil. Exot. t. 49.

L. procera, Spreng. Southern Hemisphere. Hook. Gard. Ferns, t. 53.

L. punctulata, Kze. South Africa. *Scolopendrium Krebsii*, Kze; Bot. Mag. t. 4768.

var. *blechnoidea*, Baker. South Africa.

L. Spicant, Desv. North temperate zone. *Blechnum boreale*, Sw.; Hook. Brit. Ferns, t. 40.

var. *imbricata*, Hort.

L. tuberculata, J. Sm. *See* procera.

L. zamiaefolia, Hort. *See* boryana.

L. zumoides, Gardn. *See* boryana.

Lomariopsis. *See* Acrostichum.

L. buxifolium, Fée. *See* A. sorbifolium, var. buxifolium.

Lonchitis, L.

L. pubescens, Willd. Tropics.

var. *mollissima*, Hort. Tropical Africa. *Compteris brazziana*, Linden.

Lopholepis. *See* Polypodium.**Lophosoria. *See* Alsophila.**

Lorinseria. *See* Woodwardia.

Lotzea. *See* Asplenium.

Loxoscaphe. *See* Davallia.

Lygodictyon. *See* Lygodium.

L. Fosteri, J. Sm. *See* Lygodium reticulatum.

Lygodium, Sw.

L. circinatum, Sw. *See* dichotomum.

L. dichotomum, Sw. Tropical Asia. Hook. & Grev. Ic. Fil. t. 55.

L. japonicum, Sw. Tropics of the Old World.

L. palmatum, Sw. United States. Hook. Fil. Exot. t. 24.

L. reticulatum, Schkuhr. Polynesia, East Tropical Australia.

L. scandens, Sw. Tropics of the Old World.

var. *microphyllum*, (*R. Br.*) Baker. Australia.

L. volubile, Sw. West Indies, Tropical America.

Marattia, J. E. Sm.

M. alata, J. E. Sm. West Indies, Central America.

M. attenuata, Labill. New Caledonia. Ill. Hort. 1876. t. 246.

M. Ascensionis, J. Sm. *See* fraxinea, var. purpurascens.

M. Burkei, Baker. Colombia. Gard. Chron. 1897, xxii. 425, fig. 129.

M. Cooperi, Hort. *See* attenuata.

M. cristata, J. Sm. *See* fraxinea, var. purpurascens.

Marattia, J. E. Sm.—cont.

M. elegans, Endl. *See fraxinea.*

M. fraxinea, J. E. Sm. Tropics and sub-tropics of the Old World. Bedd. Handb. Ferns Ind. 461.

var. *purpurascens*, (*De Vriese*) Baker. Ascension Island.

M. Kaulfussii, J. Sm. West Indies, Tropical America.

M. laxa, Kze. Mexico.

M. Moorei, Hort. *See attenuata.*

M. sorbifolia, Bory. *See fraxinea.*

Meniscium, Schreb.

M. deltigerum, Wall. *See Acrostichum virens.*

M. reticulatum, Schkuhr. West Indies, Tropical America
var. *macrophyllum*, Baker. Brazil.

M. serratum, Cav. West Indies, Tropical America.

M. simplex, Hook. Tropical and sub-tropical Asia. Hook.
Fil. Exot. t. 73.

Mertensia. *See Gleichenia.*

Mesochlæna. *See Didymochlæna.*

Microlepia. *See Davallia.*

Microsorium. *See Polypodium.*

Mohria, Sw.

M. Caffrorum, Desv. South Africa.

var. *achilleæfolia*, Hort.

M. thurifraga, Sw *See Caffrorum.*

Myriopteris. *See Cheilanthes.*

Neottopteris. *See Asplenium.*

Nephrodium, L. C. Rich.

N. acrostichoides, Michx. *See Aspidium.*

N. affine, Lowe. *See Felix-Mas.*

N. albopunctatum, Desv. Tropical and South Africa,
Mascarene Islands, Fiji. Hook. Fil. Exot. t. 89.

N. areolatum, Hort. *See Woodwardia areolata.*

N. atratum, Wall. *See hirtipes.*

N. Blumei, Hook. *See intermedium.*

N. canum, Baker. Himalaya.

N. chrysolobum, Féé. Brazil and Colombia.

N. cicutarium, Baker. Tropics, ubiquitous. *Aspidium coadunatum*, Hook. & Grev. Ic. Fil. t. 202.

N. cochleatum, D. Don. *See Felix-mas, var.*

N. confluens, F. Muell. Queensland.

N. conterminum, Desv. West Indies, Tropical America.

N. coruscum, Hort. *See Aspidium varium.*

N. crinitum, Desv. Mascarene Islands. *Polypodium crinitum*, Hook. & Grev. Ic. Fil. t. 66.

N. cristatum, Michx. Europe, North America. Hook.
Brit. Ferns, t. 17.

var. *clintonianum*, Hort.


var. *uliginosum*, Baker.

N. cuspidatum, Baker. Northern India, Yunnan.
Lastrea cuspidata, Bedd. Handb. Ferns Ind. 232.

Nephrodium, L. C. Rich.—cont.

- N. decompositum, R. Br.** Australia, New Zealand, Polynesia.
 var. *glabellum*, *Baker.* Australia, New Zealand.
 var. *Shepherdii*, (*Kze*) *Baker.* Australia, New Zealand
- N. decurrens, Baker.** Tropical Asia.
 var. *Thwaitesii*, *J. Sm.* Ceylon.
- N. decursivopinnatum, Baker.** China and Japan.
- N. dilatatum, Desv.** See *spinulosum*, var.
- N. dissectum, Desv.** Tropics of the Old World.
 var. *membranifolium*, (*Presl*) *Baker.* Malaya.
- N. effusum, Baker.** West Indies, Tropical America.
 var. *divergens*, *Baker.*
- N. erythrosorum, Hook.** China, Japan. Hook. Sp. Fil. iv. t. 253.
- N.* *Filix-mas, L. C. Rich.*** Temperate regions, ubiquitous. Hook. Brit. Ferns, t. 15.
 var. *cochleatum, Hook.* India.
- N. fragrans, Richardson.** North Asia, North America.
- N. glabellum, A. Cunn.** See *decompositum*.
- N. goldieanum, Hook.** North America. Hook. & Grev. Ic. Fil. t. 102.
- N. Grisebachii, Baker.** Cuba.
- N. hirtipes, Hook.** Tropical Asia. *Lastrea hirtipes*, Bedd. Handb. Ferns Ind. 233.
- N. hispidum, Hook.** Australia, New Zealand.
- N. intermedium, Baker.** Tropical Asia.
- N. Jenmani, Baker.** Jamaica.

* For garden varieties, see p. 175.



Nephrodium, L. C. Rich.—cont.

- N. lacerum**, *Baker.* Japan.
- N. latifolium**, *Baker.* Polynesia.
- N. lepidum**, *Moore.* Polynesia.
- N. leuzeanum**, *Hook.* Tropical Asia, Polynesia. *Pleocnemia leuzeana*, Presl; Gard. Chron. 1874, ii. 355, fig. 74.
- N. macrophyllum**, *Baker.* West Indies Tropical America.
var. *calvatum*, *Hort.*
- N. marginale**, *Michx.* North America.
- N. Maximowiczii**, *Baker.* China.
- N. membranifolium*, Presl. See *dissectum*.
- N. molle**, *Desv.* Ubiquitous in warm countries.
var. *polydactylon*, *Hort.* Tropics.
- N. montanum**, *Baker.* Europe, North America.
N. Oreopteris, Desv.; Hook. Brit. Ferns, t. 14.
- N. nevadense**, *Baker.* California.
- N. noveboracense**, *Desv.* North America.
- N. Oreopteris*, Desv. See *montanum*.
- N. patens**, *Desv.* Tropics and slightly beyond.
- N. patulum**, *Baker.* West Indies, Tropical America.
- N. polymorphum**, *Baker.* India, Malaya.
- N. prolificum*, Maxim. See *erythrosorum*.
- N. recedens**, *Hook.* Philippines, India, Ceylon.
- N. Richardsi**, *Baker.* New Caledonia.
var. *multifidum*, *Hort.*

Nephrodium, L. C. Rich.—cont.

- N. rigidum**, *Desv.* North temperate zone. *Hook. Brit. Ferns*, t. 16.
 var. *americanum*, *Hook.* North America.
- N. sanctum**, *Baker.* West Indies, Guatemala.
- N. scabrosum**, *Baker.* Nilgherries. *Lastrea scabrosa*, *Bedd. Hand. Ferns Ind.* 263.
- N. setigerum**, *Baker.* Japan, Tropical Asia, Polynesia.
 var. *cristatum*, *Hort.*
- N. Sieboldii**, *Hook.* *Fil. Exot.* t. 31. Japan.
- N. Sloanei**, *Baker.* West Indies, Tropical America.
- N. *spinulosum**, *Desv.* North temperate zone.
 var. *Boottii*, *Baker.* United States.
 var. *dilatatum*, *Hook.* Europe. *Hook. Brit. Ferns*, t. 19.
- N. subbiauritum**, *Hook.* Madagascar.
- N. subtriphyllum**, *Baker.* Tropical Asia, Polynesia, Mascarene Is.
- N. tenericaule*, *Hook.* See *setigerum*.
- N. Thelypteris**, *Desv.* North and South temperate zones.
Hook. Brit. Ferns, t. 13.
- N. trichophorum**, *Baker.* Guadeloupe.
- N. truncatum**, *Presl.* India, Malaya, Polynesia, Australia.
- N. unitum**, *R. Br.* Tropical and subtropical regions.
- N. variolosum**, *Baker.* India, Malaya.
- N. villosum**, *Presl.* West Indies, Tropical America.

Nephrolepis, Schott.

- N. acuta**, *Presl.* Tropics, ubiquitous.

* For garden varieties. see p. 179.

Nephrolepis, Schott—cont.

- N. Bausei**, *Hort.* Garden origin.
- N. biserrata**, Schott. *See acuta.*
- N. bostoniensis**, *Hort.* Garden origin.
- N. cordifolia**, *Baker.* Tropics, ubiquitous. *Bedd. Handb. Ferns Ind.* 283.
- var. *compacta*, *Hort.*
 - var. *crispato-congesta*, *Hort.*
 - var. *edmontonensis*, *Hort.*
 - var. *pectinata*, *Baker.* Tropical America.
 - var. *philippinensis*, *Hort.* Philippines.
- N. davallioides**, *Kze.* Java. *Hook. Fil. Exot. t.* 60.
- var. *furcans*, *Hort.*
 - var. *multiceps*, *Hort.*
- N. Duffii**, *Moore.* Australia. *Gard. Chron.* 1878, ix. 623, fig. 113.
- N. ensifolia**, Presl. *See acuta.*
- N. exaltata**, *Schott.* Tropics, ubiquitous. *Bedd. Ferns S. Ind. t.* 93.
- var. *Barteri*, *Hort.* West Tropical Africa.
 - var. *hirsutula*, *Baker.* Malaya, Tropical Australia.
- N. Faulkneri**, *Hort.* Garden origin.
- N. floccigera**, *Hort.* Garden origin.
- N. Fosteri**, *Hort.* Garden origin.
- N. Hestoni**, *Hort.* Garden origin.
- N. pectinata**, Schott. *See cordifolia*, var.
- N. philippinensis**, *Hort.* *See cordifolia*, var.
- N. Pluma**, *Moore.* Madagascar. *Gard. Chron.* 1878, ix. 589, fig. 108.

Nephrolepis, Schott—cont.

N. ramosa, *Moore*. Tropics of the Old World.

N. recurvata, *Hort.* *See exaltata*.

N. rufescens, *Presl*. Tropics.

var. **Mayi**, *Hort.*

var. **tripinnatifida**, *Baker*. Solomon Islands.

N. tuberosa, *Presl*. *See cordifolia*.

N. undulata, *J. Sm.* *See cordifolia*.

N. Westoni, *Hort.* Garden origin.

N. Wittboldi, *Hort.* Garden origin.

Neurocallis. *See Acrostichum*.

Niphobolus. *See Polypodium*.

Niphopsis. *See Polypodium*.

Nothochlæna. *See Notholæna*.

Notholæna, R. Br.

N. distans, *R. Br.* Australia, New Zealand, New Caledonia.

N. eckloniana, *Kze.* South Africa.

N. ferruginea, *Hook.* West Indies, Tropical America.

N. flavens, *Moore*. Central America. *Gymnogramme flavens*, *Sw.*; *Hook. Fil. Exot. t. 47*.

N. Hookeri, *Lowe*. *See nivea*.

N. laevis, *Martens & Gal.* *See sinuata*.

N. lanuginosa, *Desv.* S. Europe, Atlantic Islands, Australia.

N. Marantæ, *R. Br.* S. Europe, N. Africa, temp. Asia.

Notholæna, R. Br.—cont.

N. rufa, Presl. *See ferruginea.*

N. sinuata, Kaulf. Tropical America.

N. trichomanoides, R. Br. Jamaica, Cuba.

Odontosoria. *See Davallia.***Oleandra, Cav.**

O. articulata, Cav. Tropical and S. Africa, Mascarene Islands.

O. nodosa, Presl. West Indies, Tropical America.

Olfersia. *See Acrostichum.***Onoclea, L.**

O. germanica, Willd. North temperate zone.

O. pennsylvanica, Willd. *See germanica.*

O. sensibilis, L. Northern Asia, North America.

Onychium, Kaulf.

O. japonicum, Kze. Japan, Himalaya.

O. lucidum, Spreng. *See japonicum.*

Ophioglossum, L.

O. vulgatum, L. Ubiquitous in cold and temperate regions. Hook. Brit. Ferns, t. 46.

Osmunda, L.

O. bipinnata, Hook. Fil. Exot. t. 9. Hong Kong.

O. cinnamomea, L. North and Central America, West Indies, Northern Asia. Hook. Gard. Ferns, t. 45.

Osmunda, L.—cont.

- O. claytoniana, L.** North America, Himalaya.
- O. interrupta, Michx.** See *claytoniana*.
- O. japonica, Thunb.** See *regalis*.
- O. javanica, Bl.** From Kamschatka to Java.
- O. *regalis, L.** North and South temp. regions. Hook. Brit. Ferns, t. 45.
var. *japonica, Milde.* Japan.
- O. spectabilis, Willd.** See *regalis*.

Pæsia. See *Pteris*.

Paragamma. See *Polypodium*.

Parkeria. See *Ceratopteris*.

Pellæa, Link.

- P. adiantifolia, Hort.** See *hastata*.
- P. atropurpurea, Link.** North and Central America.
Lowe, Ferns, iii. t. 30.
- P. cordata, J. Sm.** Mexico, Western South America.
Allosorus cordatus, Presl; Bot. Mag. t. 4698.
var. *flexuosa, Baker.* Western South America.
Allosorus flexuosus, Kaulf.; Bot. Mag. t. 4762.
- P. densa, Hook.** North America.
- P. falcata, Fée.** Tropical Asia and Australia.
- P. geraniifolia, Fée.** Tropics. *Pteris geraniifolia, Raddi*; Hook. Ic. Pl. t. 915.
- P. gracilis, Hook.** Northern Asia, North America.
- P. hastata, Link.** Tropical and South Africa. Hook. Fil. Exot. t. 50.

* For garden varieties, see p. 179.

Pellæa, Link—cont.

- P. intramarginalis, J. Sm.* Central America.
P. involuta, Baker. East Tropical Africa.
P. mucronata, Eaton. California, New Mexico.
P. rotundifolia, Hook. New Zealand, Norfolk Island.
 Hook. Fil. Exot. t. 48.
P. viridis, Forsk. *See hastata.*
P. wrightiana, Hook. *See mucronata.*

Phegopteris. *See Polypodium.*

P. trichodes, J. Sm. *See Nephrodium setigerum.*

Phlebodium. *See Polypodium.*

P. inaequale, Moore. *See Polypodium guatemalense.*

Phymatodes. *See Polypodium.*

P. vulgaris, Presl. *See Polypodium Phymatodes.*

Platycerium, Desv.

- P. æthiopicum, Hook.* Tropical Africa. Hook. Gard. Ferns, t. 9.
P. alcicorne, Desv. Australia. Bot. Reg. tt. 262-3.
 var. *Hillii, Bailey.* Queensland. *P. Hillii, Moore,* Gard. Chron. 1878, x. 428, figs. 74-5.
P. angolense, Welw. Tropical Africa.
P. biforme, Bl. Malaya. Bedd. Ferns Brit. Ind. tt. 109, 224.
P. grande, J. Sm. Tropical Asia, North Australia.
 Hook. Fil. Exot. t. 86.
P. madagascariense, Baker. Madagascar.

Platycerium, *Desv.*—*cont.*

P. Stemmaria, P. Beauv. *See* *aethiopicum*.

P. Veitchii, *Hort.* Garden origin.

P. Wallichii, *Hook.* Malay Peninsula. *Hook. Fil. Exot. t. 97.*

P. Willinckii, *Moore.* Java. *Gard. Chron. 1875*, iii. 303, fig. 56.

Platyloma. *See* *Pellaea*.

Pleocnemia. *See* *Nephrodium*.

Pleopeltis. *See* *Polypodium*.

Pleuridium. *See* *Polypodium*.

Pœcilopteris. *See* *Acrostichum*.

Polybotrya. *See* *Acrostichum*.

Polypodium, *L.*

P. adnascens, *Sw.* Tropical Asia and Africa. *Niphobolus adnascens*, Kaulf. ; *Hook. Gard. Ferns*, t. 19.

P. albopunctatum, *Linden.* *See* *crassifolium*, var.

P. albosquamatum, *Bl.* Malaya. *Hook. Gard. Ferns*, t. 47.

P. alpestre, *Hoppe.* North temperate zone. *Hook. Brit. Ferns*, t. 6.

var. *flexile*, *Moore.* Scotland.

P. americanum, *Hook.* Andes of Ecuador.

P. amphostemon, *Kze.* *See* *angustifolium*.

Polypodium, L.—cont.

P. angulatum, Willd. Malay Islands.

P. angustatum, Sw. Malay Peninsula and Islands. Australia. *Niphobolus angustatus*, Spreng.; Hook. Gard. Ferns, t. 20.

P. angustifolium, Sw. West Indies, Tropical America.

P. appendiculatum, Wall. India. Hook. Fil. Exot. t. 87.

P. areolatum, H. B. & K. See *aureum*, var.

P. attenuatum, H. B. & K. West Indies, Tropical America.

P. aureum, L. West Indies, Tropical America.

var. *areolatum*, Baker. Brazil.

var. *glaucum*, Hort.

var. *Mayi*, Hort.

var. *pulvinatum*, Baker.

var. *sporodocarpum*, (Willd.) Baker.

P. Billardieri, R. Br. Australia and New Zealand.

P. brevitolium, Link. See *Phylltidis*, var. *linnaeanum*.

P. calcareum, J. E. Sm. See *Dryopteris*, var. *robertianum*.

P. cambricum, L. See *vulgare*, var.

P. cameroonianum, Hook. Cameroon Mountains.

P. Catharinæ, J. Sm. See *loriceum*.

P. chnoodes, Spreng. West Indies, Venezuela.

P. colpodes, Kze. See *plesiosorum*.

P. conjugatum, Lam. Tropical Asia. *P. coronans*, Wall.; Hook. Fil. Exot. 91.

P. contiguum, Wall. See *longifolium*.

Polypodium, L.—*cont.*

P. coronans, Wall. *See conjugatum.*

P. crassifolium, L. West Indies, Tropical America.
var. *albopunctatissimum*, (*J. Sm.*) Baker. Brazil.

P. cuspidatum, Bl. *See persicæfolium.*

P. decurrens, Raddi. Tropical America.

P. dissimile, Hook. *See chnoodes.*

P. distans, Raddi. *See fraxinifolium.*

P. divergens, Willd. *See Nephrodium effusum, var.*

P. diversifolium, Sw. Tropical America. Hook. Gard. Ferns, t. 5.

P. Dryopteris, L. North temperate zone. Hook. Brit. Ferns, t. 4.

var. *robertianum*, Ruprecht. *P. robertianum*, Hoffm.; Hook. Brit. Ferns, t. 5.

P. effusum, Sw. *See Nephrodium.*

P. elasticum, L. C. Rich. Tropical America.

P. ensifolium, Willd. *See angustifolium.*

P. falcatum, Kellogg. California.

P. Fendleri, Eaton. Venezuela, Colombia.

P. fossum, Baker. Garden origin.

P. fraxinifolium, Jacq. Tropical America. Jacq. Ic. t. 639.

P. glaucophyllum, Kze. West Indies, Tropical America. West Africa. *Goniophlebium glaucophyllum*, J. Sm.; Gard. Chron. 1874, ii. 7, fig. 1.

var. *glabrum*, Moore.

var. *viride*, Hort.

P. grande, Presl. *See macropterum, var. splendidum.*

Polypodium, L.—cont.

- P. guatemalense**, Hook. Central America.
- P. harpeodes**, Link. See *loriceum*.
- P. hastatum**, Thunb. Japan.
- P. hemionitideum**, Wall. North India, South China.
- P. Heracleum**, Kze. Java, Philippines. Hook. Gard. Ferns, t. 1.
- P. hexagonopterum**, Michx. North America. Hook. & Grev. Ic. Fil. t. 210.
- P. hirsutum**, Sw. Brazil.
- P. hydrophilum**, Kze. Dominica.
- P. incanum**, Sw. Tropical and South Africa, America from Ohio southwards.
- P. incurvatum**, Bl. Malay Peninsula.
- P. irioides**, Lam. Tropics and subtropics of the Old World. Hook. Fil. Exot. t. 4.
var. **ramocristatum**, Hort.
- P. juglandifolium**, D. Don. Himalaya.
var. **tenuicauda**, Baker.
- P. kariwinskianum**, A. Br. See *plebeium*.
- P. Knightiæ**, Hort. Australia. Gard. Chron. 1903, xxxiii. 245, fig. 99.
- P. lachnopus**, Wall. Northern India.
- P. lævigatum**, Cav. Tropical America.
- P. lanceolatum**, L. Tropics.
- P. lapathifolium**, Poir. See *lævigatum*.
- P. leiorhizon**, Wall. India, China. Hook. Fil. Exot. t. 25.

Polypodium, L.—*cont.*

- P. lepidopteris*, Kze. Tropical America.
 var. *sepultum*, Baker. Tropical America.
- P. lineare*, Thunb. Tropical and subtropical Asia,
 Tropical and subtropical Africa.
- P. Lingua*, Sw. Manchuria to Malaya.
 var. *corymbiferum*, Hort.
 var. *Heteractis*, Baker. Himalaya.
- P. longifolium*, Mett. Malaya. *P. contiguum*, Wall. ;
 Hook. Fil. Exot. t. 20.
- P. longipes*, Link. See *Phymatodes*, var.
- P. longissimum*, Bl. Tropical Asia.
- P. loriceum*, L. West Indies, Tropical America.
 var. *hirsutum*, Hort.
 var. *latipes*, Baker. Brazil.
 var. *pinnatifidum*, Baker.
- P. lycopodioides*, L. Tropics, ubiquitous.
 var. *maximum*, Hort.
 var. *owariense*, (Desv.) Baker. Tropical Africa.
 var. *salicifolium*, Hook. Tropical America.
 var. *squamulosum*, Baker. South Brazil.
- P. macropterum*, Kaulf. Brazil.
 var. *splendidum*, Baker. South Brazil.
- P. macrourum*, Baker. Australia.
- P. menisciifolium*, Langsd. & Fisch. See *neriifolium*.
- P. meyenianum*, Schott. Philippines.
- P. multilineatum*, Wall. Northern India.
- P. musæfolium*, Bl. Malaya.
- P. neriifolium*, Schkuhr. West Indies, Tropical America.
 var. *albopunctatum*, (Raddi). Tropical America.

Polypodium, L.—cont.

- P. nigrescens**, Bl. India, Malaya, Polynesia. Hook. Fil. Exot. t. 22.
 var. *grandiceps*, Hort.
- P. ovariense**, Desv. See *lycopodioides*.
- P. palmatum**, Bl. Malaya.
- P. pectinatum**, L. West Indies, Tropical America. Hook. Gard. Ferns, t. 10.
 var. *Paradisæ*, Baker. Brazil.
 var. *plumosum*, Hort.
- P. peltideum**, Link. See *Phymatodes*.
- P. pendulum**, Sw. West Indies, Tropical America.
- P. pennigerum**, Forst. New Zealand.
- P. percussum**, Cav. Tropical America. Hook. Fil. Exot. t. 59.
- P. persicæfolium**, Desv. Malaya.
 var. *grandidens*, Kze. Java.
- P. *Phegopteris**, L. North temperate zone. Hook. Brit. Ferns, t. 3.
- P. Phyllitidis**, L. West Indies, Tropical America.
 var. *linnæanum*, Hook.
 var. *obscurum*, (Mett.) Baker.
- P. Phymatodes**, L. Tropical Asia and Africa.
 var. *cristatum*, Hort.
 var. *longipes*, (Link) Baker.
 var. *terminale*, (Link) Baker.
- P. pictum**, Baker. Polynesia.
- P. piloselloides**, L. West Indies, Tropical America. Hook. Gard. Ferns, t. 18.

* For garden varieties, see p. 181.

Polypodium, L.—cont.

- P. platyphyllum*, *Hook.* West Indies, Tropical America.
- P. plebeium*, *Schlecht.* Central and South America.
Hook. *Gard. Ferns*, t. 48.
var. *madrense*, (*J. Sm.*). Central America.
- P. plesiosorum*, *Kze.* Mexico to Venezuela.
- P. Plumula*, H. B. & K. *See elasticum.*
- P. preslianum*, Link. *See neriifolium.*
- P. propinquum*, *Wall.* China, India, Java.
var. *Laurentii*, *Christ.* Tropical Africa. De Wildem.
Etudes Fl. Bas-Congo, 6, t. 2.
- P. pulvinatum*, Link. *See aureum*, var.
- P. pustulatum*, *Forst.* Australia and New Zealand.
- P. quercifolium*, *L.* Tropical Asia, Queensland.
- P. repens*, *L.* West Indies, Tropical America.
var. *undulatum*, *Hort.*
- P. reptans*, *Sw.* West Indies, Tropical America.
- P. rigidulum*, *Sw.* Malaya, Fiji, Queensland.
- P. robertianum*, Hoffm. *See Dryopteris*, var.
- P. rostratum*, *Hook.* Northern India.
- P. rupestre*, *Bl.* Malaya.
- P. Schneideri*, *Hort.* Garden hybrid. *Journ. Hort.*
ser. 3, xxix. 85, fig. 13.
- P. Scouleri*, *Hook.* & *Grev.* *Ic. Fil.* t. 56. N. W.
America.
- P. sepultum*, Kaulf. *See lepidopteris*, var.
- P. serrulatum*, Mett. Tropical Africa and America.
Xiphopteris serrulata, Kaulf. ; *Hook. Gard. Ferns.*
t. 44.

Polypodium, L.—cont.

- P. simplex*, Sw. *See lineare.*
- P. sparsiflorum**, Hook. West Tropical Africa.
- P. spherocephalum*, Wall. *See angustatum.*
- P. sporodocarpum*, Willd. *See aureum*, var.
- P. squamulosum*, Kaulf. *See lycopodioides.*
- P. stigmaticum**, Presl. Colombia.
- P. stigmosum**, Sw. India, Malaya.
- P. subauriculatum**, Bl. India, Malaya, Polynesia, Queensland.
- P. superficiale**, Bl. China, India.
- P. tenuiosum*, Hort. *See angustifolium.*
- P. taxifolium**, L. West Indies, Tropical America.
- P. tenuicauda*, Hook. *See juglandifolium*, var.
- P. terminale*, Link. *See Phymatodes*, var.
- P. tetragonum**, Sw. West Indies, Tropical America.
- P. tricuspe**, Sw. Japan.
- P. vacciniifolium**, Langsd. & Fisch. West Indies, Tropical America.
var. **album**, Hort.
- P. vacillans*, Link. *See loriceum.*
- P. venosum*, Desv. *See stigmaticum.*
- P. verrucosum**, Wall. Malaya. Hook. Gard. Ferns, t. 41.
- P. *vulgare**, L. North temperate zone. Hook. Brit. Ferns, t. 2.
- P. Willdenovii**, Bory. Tropical Africa, Mascarene Islands. Hook. Gard. Ferns, t. 35.

* For garden varieties, *see p. 181.*

Polypodium, L.—cont.

P. Xiphias, Baker. Southern Polynesia.

Polystichum. *See Aspidium.*

Prosaptia. *See Davallia.*

Pseudathyrium. *See Polypodium.*

Pteridella. *See Pellæa.*

P. viridis, Mett. *See Pellæa hastata.*

Pteris, L.

P. aquilina, L. Cosmopolitan. Hook. Brit. Ferns, t. 38.

P. arguta, Ait. South Europe, Azores &c.

P. albolineata, Hort. *See cretica*, var.

P. argyræa, Moore. *See quadriaurita*, var.

P. Bausei, Hort. *See semipinnata*, var.

P. biaurita, L. Tropics. Hook. & Grev. Ic. Fil. t. 142.

var. **argentea**, Hort.

var. **nemoralis variegata**, Hort.

P. Childsii, Hort. Garden origin.

P. crenata, Sw. *See ensiformis.*

P. cretica, L. Tropical and subtropical regions.

var. **albolineata**, Hook. Bot. Mag. t. 5194.

var. **Alexandrae**, Hort.

var. **crispata**, Hort.

var. **Maissonieri**, Hort.

var. **major**, Hort.

var. **Mayi**, Hort.

var. **nobilis**, Hort.


var. **Wimsetti**, Hort.

Pteris, L.—*cont.*

- P. denticulata*, Sw. West Indies, Tropical America.
Hook. & Grev. Ic. Fil. t. 28.
- P. Drinkwateri*, Hort. *See pellucida.*
- P. droogmansiana*, De Wild. Belgian Congo.
- P. Duvalii*, (*Bellair*). Garden hybrid. *Dryopteris Duvalii*, Bellair. Rev. Hort. 1897, 563, fig. 168.
- P. elata*, Ag. Tropical America.
var. *karsteniana*, Hort.
- P. ensiformis*, Burm. Tropical Asia and Australia, Polynesia.
var. *cristata*, Hort.
var. *Reginæ*, Hort.
var. *Victoriæ*, Hort.
- P. felosma*, J. Sm. *See quadriaurita.*
- P. flabellata*, Thunb. Tropical and South Africa.
- P. geraniifolia*, Raddi. *See Pellæa.*
- P. Ghiesbreghtii*, J. Sm. *See laciniata.*
- P. grevilleana*, Wall. Northern India.
var. *variegata*, Hort.
- P. heterophylla*, L. West Indies. Bot. Mag. t. 4925.
var. *internata*, Baker. West Indies.
- P. incisa*, Thunb. Tropical and subtropical regions.
var. *Vespertilionis*, (Labill.). Australia.
- P. internata*, Moore. *See heterophylla*, var.
- P. laciniata*, Willd. West Indies, Tropical America.
var. *Ghiesbreghtii*, Baker. Mexico.
- P. leptophylla*, Sw. Brazil. Hook. Gard. Ferns, t. 23.
var. *argentea*, Hort.

Pteris, L.—cont.

- P. longifolia, L.* Tropical and subtropical regions.
var. *Mariesii, Hort.* Japan.
- P. ludens, Wall.* India, Malaya. *Doryopteris ludens,* Bedd. Handb. Ferns Ind. 119.
- P. Maissonieri, Hort.* See *cretica*, var.
- P. moluccana, Bl.* Malaya.
- P. nemoralis, Willd.* See *biaurita*, var.
- P. nobilis, Hort.* See *palmata*, var.
- P. palmata, Willd.* West Indies, Tropical America.
Hook. Gard. Ferns, t. 22.
var. *nobilis, Hort.*
- P. pedata, L.* West Indies, Tropical America. Hook.
Fil. Exot. t. 34.
- P. pellucida, Presl.* India, Malaya, Tropical Africa.
- P. pulchra, Schl.* Mexico.
- P. pungens, Willd.* West Indies.
- P. quadriaurita, Retz.* Tropics, ubiquitous. Bot. Mag.
t. 5183.
var. *argyræa, (Moore) Baker.*
var. *rubicaulis, (J. Sm.) Baker.*
var. *tricolor, Hort.*
- P. sagittifolia, Raddi.* Brazil. Hook. Fil. Exot. t. 39.
- P. scaberula, A. Rich.* New Zealand.
- P. semipinnata, L.* Tropical and Sub-tropical Asia.
Hook. Gard. Ferns, t. 59.
var. *Bausei, Hort.*
var. *gigantea, Hort.*



Pteris, L.—cont.

P. serrulata, *L. fil.* China, Japan, Natal.

var. **cristata**, *Hort.*

var. **voluta**, *Hort.*

P. straminea, *Mett.* Chili.

P. sulcata, *Link.* See *quadriaurita*.

P. tremula, *R. Br.* Australia and New Zealand.

P. Vespertilionis, *Labill.* See *incisa*, var.

P. Victoriae, *Hort.* See *ensiformis*, var.

P. viridis, *Forsk.* See *Pellaea hastata*.

P. wallichiana, *Ag.* Himalaya, Japan, Philippines.

Pteropsis. See *Tænitis*.

Pycnopteris. See *Nephrodium*.

Rhipidopteris. See *Acrostichum*.

Sagenia. See *Nephrodium*.

Salpichlæna. See *Blechnum*.

Schellolepis. See *Polypodium*.

Schizocæna. See *Cyathea*.

Schizoloma. See *Lindsaya*.

Scolopendrium, *J. E. Sm.*

S. brasiliense, *Kze.* Brazil.

S. Delavayi, *Franch.* China, Munipur. Journ. Linn. Soc. xxv. t. 41.

Scolopendrium, J. E. Sm.—cont.

S. Krebsii, Kze. *See Lomaria punctulata*.

S. rhizophyllum, Hook. North America. *Camptosorus rhizophyllus*, Presl; Hook. Fil. Exot. t. 85.

S.*vulgare, J. E. Sm. North temperate zone. Hook. Brit. Ferns, t. 37.

Selliguea. *See Gymnogramme*.

Sitolobium. *See Dicksonia*.

Soromanes. *See Acrostichum Cænopteris*.

Stegania. *See Lomaria*.

Stenochlæna. *See Acrostichum*.

Stenoloma. *See Davallia*.

Stenosemia. *See Acrostichum*.

Struthiopteris. *See Onoclea*.

Tæniopsis. *See Vittaria*.

Thyrsopteris, Kze.

T. elegans, Kze. Juan Fernandez.

Todea, Willd.

T. africana, Willd. *See barbara*.

T. barbara, Moore. South Africa, Australia, New Zealand. Bot. Mag. t. 5954.

* For garden varieties, see p. 181.

Todea, Willd.—cont.

- T. Fraseri**, Hook. & Grev. Ic. Fil. t. 101. Australia,
New Caledonia.
- var. *wilkesiana*, Baker. Fiji, New Hebrides, New
Guinea. *T. wilkesiana*, Brackenr.; Ill. Hort. 1875,
t. 226.
- T. grandipinnula*, Moore. See *Moorei*.
- T. hymenophylloides**, Rich. & Less. New Zealand.
Hook. Gard. Ferns, t. 54.
- T. intermedia*, Hort. See *hymenophylloides*.
- T. Moorei**, Baker. Lord Howe Island.
- T. pellucida*, Carm. See *hymenophylloides*.
- T. rivularis*, Sieb. See *barbara*.
- T. superba**, Col. New Zealand. Ill. Hort. 1870, t. 23.
- T. wilkesiana*, Brackenr. See *Fraseri*, var.

Trichomanes, L.

- T. alatum**, Sw. West Indies, Tropical America.
Hook. & Grev. Ic. Fil. t. 11.
- var. *attenuatum*, Griseb. West Indies.
- T. angustatum*, Carm. See *tenerum*.
- T. attenuatum*, Hook. See *alatum*, var.
- T. auriculatum**, Bl. Tropical Asia.
- T. Bancroftii**, Hook. & Grev. West Indies, Tropical
America. Hook. Gard. Ferns, t. 56.
- T. bipunctatum*, Poir. See *Filicula*.
- T. Bojeri*, Hook. & Grev. See *cuspidatum*.
- T. boschianum*, Sturm. See *radicans*, var.
- T. Colensoi**, Hook. f. New Zealand. Hook. Ic. Pl.
t. 979.

Trichomanes, L.—cont.

- T. crispum**, L. West Indies, Tropical America. Hook.
Gard. Ferns, t. 27.
- T. cuspidatum**, Willd. Mascarene Islands.
- T. elongatum**, Cunn. See rigidum, var.
- T. exsectum**, Kze. Chili, Juan Fernandez.
- T. Filicula**, Bory. Tropical Asia and Africa.
- T. fœniculaceum**, Bory. Mascarene Is., Borneo,
Australia.
- T. hirtellum**, Poir. See Hymenophyllum.
- T. humile**, Forst. Java, New Zealand, Polynesia. Hook.
& Grev. Ic. Fil. t. 85.
- T. Kaulfussii**, Hook & Grev. West Indies, Tropical
America.
- T. labiatum**, Jenm. British Guiana.
- T. luschnathianum**, Presl. See radicans, var.
- T. Lyallii**, Hook. New Zealand.
- T. maximum**, Bl. Malaya, Polynesia, N. Australia.
- T. membranaceum**, L. West Indies, Tropical America.
- T. muscoides**, Sw. Tropics.
- T. parviflorum**, Poir. See fœniculaceum.
- T. parvulum**, Poir. Tropics of the Old World.
- T. pinnatum**, Sw. West Indies, Tropical America. Hook.
Gard. Ferns, t. 8.
- T. Prieurii**, Kze. West Indies, Tropical America.
- T. pyxidiferum**, L. Tropics, ubiquitous. Hook. & Grev.
Ic. Fil. t. 206.
- T. radicans**, Sw. Tropical, subtropical and temperate
regions.
- var. **Andrewsii**, Newm. Ireland.

Trichomanes, L.—cont.**T. radicans, Sw.—cont.**

var. *boschianum*, (*Sturm*) *Baker*. Alabama.

var. *luschnathianum*, *Baker*. Brazil.

var. *luschnathianum prolongatum*, *Hort.*

var. *proliferum*, *Hort.*

T. reniforme, Forst. New Zealand. Hook. Fil. Exot. t. 2.

T. reptans, Sw. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 32.

T. rigidum, Sw. Tropics, ubiquitous.

var. *elongatum*, *Baker*. New Zealand.

T. scandens, L. West Indies, Central America.

T. sinuosum, L. C. Rich. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 13.

T. spicatum, Hedw. West Indies, Tropical America. Hook. Gard. Ferns, t. 60.

T. tenerum, Spreng. Tropical America.

T. trichoideum, Sw. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 199.

T. venosum, R. Br. Australia and New Zealand.

Vittaria, J. E. Sm.

V. lineata, Sw. Tropical and subtropical regions.

Woodsia, R. Br.

W. ilvensis, R. Br. Arctic and North temperate zone. Hook. Brit. Ferns, t. 8.

W. oregana, D. C. Eaton. North America.

W. scopolina, D. C. Eaton. North America.

Woodwardia, J. E. Sm.

W. angustifolia, J. E. Sm. *See* areolata.

W. areolata, *Moore*. United States. Hook. Gard. Ferns, t. 61.

W. Harlandii, *Hook.* Hong Kong. Hook. Fii. Exot. t. 7.

W. japonica, *Sw.* China and Japan.

W. radicans, *Sw.* North temperate zone.

var. *americana*, *Hook.* Southern United States.

var. *Brownii*, *Hort.*

var. *cristata*, *Hort.* *See* var. *Brownii*.

var. *plumosa*, *Hort.*

W. virginica, *J. E. Sm.* North America.

Xiphopteris. *See* Polypodium.

II.—FERN ALLIES.

Azolla, Lam.

- A. caroliniana**, Willd. Tropical and subtropical America.
A. pinnata, R. Br. Australia.

Bernhardia. See *Psilotum*.

Equisetum, L.

- E. arvense**, L. North temperate and Arctic regions.
 Hook. Brit. Ferns, t. 60.
E. fluviatile, L. See *limosum*.
E. hyemale, L. North temperate and Arctic regions.
 Hook. Brit. Ferns, t. 64.
E. limosum, L. North temperate and Arctic regions.
 Hook. Brit. Ferns, t. 62.
E. maximum, Lam. North temperate regions.
E. ramosissimum, Desf. Warm temperate and tropical regions.
E. robustum, A. Br. North and central America.
E. scirpoides, Michx. North temperate and Arctic regions.

Isoetes, L.

- I. lacustris**, L. Europe and North America. Hook.
 Brit. Ferns, t. 55.
I. malivernianum, Cesati & De Not. Italy.

Lycopodium, L.

- L. carinatum**, Desv. India, Malaya, Polynesia. *L. acrostachyum*, Wall.; Hook. & Grev. Ic. Fil. t. 181.
- L. clavatum**, L. Arctic and Alpine zones; on mountains in the tropics.
- L. dalhousieanum**, Spring. Malay Peninsula, Borneo.
- L. dichotomum**, Jacq. West Indies, Tropical America.
- L. fontinaloides**, Spring. Brazil.
- L. Hippuris**, Desv. See *squarrosum*.
- L. Hookeri**, Wall. See *squarrosum*.
- L. lucidulum**, Michx. Himalaya, China, Japan, North America.
- L. obscurum**, L. North Asia, North America.
- L. Phlegmaria**, L. Tropics of the Old World.
- L. scariosum**, Forst. New Zealand.
- L. squarrosum**, Forst. India, Malaya, Polynesia. Mascarene Is. *L. Hookeri*, Wall.; Hook. & Grev. Ic. Fil. t. 185.
- L. taxifolium**, Sw. West Indies, Tropical America. Hook. & Grev. Ic. Fil. t. 131.

Marsilea, L.

M. macra, A. Br. Australia.

M. Nardu, A. Br. Australia.

Pilularia, L.

P. globulifera, L. Europe. Hook. Brit. Ferns, t. 57.

Psilotum, Sw.

P. triquetrum, Sw. Tropics, extending to New Zealand, Japan, and Florida. Hook. Fil. Exot. t. 63.
var. mascarenica, Hort.

Salvinia, Schreb.

S. natans, Hoffm. Subtropics of the Old World.

Selaginella, Spring.

S. africana, A. Br. See Vogelii.

S. albonitens, Spring. West Indies.

S. albospica, Hort. See stenophylla, var.

S. amœna, Hort. See caulescens.

S. apus, Spring. North America.

var. elegans, Hort.

S. argentea, Spring. See caulescens, var.

S. atroviridis, Spring. Tropical Asia. *Lycopodium atroviride*, Wall. ; Hook. & Grev. Ic. Fil. t. 39.

S. bakeriana, Bailey. Queensland.

S. bellula, Moore. See inaequalifolia, var. perelegans.

S. Braunii, Baker. China.

S. cæsia, Hort. See uncinata.

S. canaliculata, Baker. Tropical Asia.

S. caudata, Spring. See canaliculata.

S. caulescens, Spring. Tropical Asia.

var. amœna, Hort.

var. argentea, Baker. Malay Peninsula.

var. japonica, MacNab. Japan.

var. minor, Hort.



Selaginella, Spring—cont.

- S. concinna**, *Spring*. Mascarene Islands.
- S. conferta**, *Moore*. *See* *canaliculata*.
- S. cuspidata**, *Link*. West Indies, Tropical America.
var. *elongata*, *Spring*. Guatemala.
- S. delicatissima**, *A. Br.* Andes.
- S. densa**, *Hort.* *See* *apus*.
- S. denticulata**, *Hort.* *See* *kraussiana*.
- S. dichroa**, *Hort.* *See* *Vogelii*.
- S. Douglasii**, *Spring*. North America.
- S. emiliana**, *Hort.*
- S. erythropus**, *Spring*. West Indies, Tropical America.
var. *minor*, *Hort.*
var. *setosa*, *Hort.*
- S. filicina**, *Spring*. *See* *hæmatodes*.
- S. flabellata**, *Spring*. Tropical and subtropical regions.
- S. flagellifera**, *Hort.* *See* *plumosa*, var.
- S. Galeottii**, *Spring*. Central America.
- S. geniculata**, *Spring*. Tropical America.
- S. grandis**, *Moore*. Borneo. *Gard. Chron.* 1882, xviii. 40, figs. 7–8.
- S. hæmatodes**, *Spring*. West Indies, Tropical America.
- S. helvetica**, *Link*. Europe, North Asia.
- S. inæqualifolia**, *Spring*. Tropical Asia.
var. *perelegans*, *Baker*. Tropical Asia. *S. bellula*,
Moore, *Gard. Chron.* 1879, xi. 173, fig. 25.
- S. involvens**, *Spring*. China, Japan, E. Himalaya.
Philippine Is.
var. *texta*, *Baker*.

Selaginella, Spring—cont.

- S. kraussiana, A. Br.** South Africa, W. Tropical Africa.
Azores.
 var. **aurea, Hort.**
 var. **Brownii, Hort.**
 var. **Stansfieldii, Hort.**
- S. lævigata, Baker.** Madagascar.
 var. **Lyallii, Baker.** Madagascar.
- S. lepidophylla, Spring.** Tropical America.
- S. leptostachya, Bailey.** See *bakeriana*.
- S. Lyallii, Spring.** See *lævigata*, var.
- S. Martensi, Spring.** Mexico. *S. pæppigiana*, Spring;
Hook. Fil. Exot. t. 56.
 var. **dichotoma, Hort.**
 var. **variegata, Hort.**
- S. Mettenii, A. Br.** Garden origin.
- S. molliceps, Spring.** Tropical Africa, Madagascar.
- S. mutabilis, Hort.** See *serpens*.
- S. nobilis, Hort.** Garden origin.
- S. patula, Spring.** West Indies.
- S. perelegans, Moore.** See *inæqualifolia*, var.
- S. pilifera, A. Br.** Texas, Central Mexico.
- S. plumosa, Baker.** Tropical Asia.
- S. Poulteri, Hort.** Azores.
- S. radiata, Baker.** Tropical America.
- S. radicata, Spring.** See *plumosa*.
- S. rubella, Moore.** Garden origin.
- S. rubricaulis, Hort.** See *suberosa*.

elaginella, *Spring*—cont.

- S. rupestris**, *Spring*. North and south temp. zones; mountains in the tropics.
- S. serpens**, *Spring*. West Indies.
- S. strobilifera**, Hort. *See rubella*.
- S. tassellata**, Hort. Brazil.
- S. texta**, Hort. *See involvens*.
- S. uncinata**, *Spring*. China.
- S. usta**, Vieill. New Caledonia.
- S. Victoriae**, Moore. Malaya, Fiji.
- S. viridangula**, *Spring*. Fiji.
- S. viticulosa**, Klotzsch. Central America, Venezuela.
- S. Vogelii**, *Spring*. Tropical Africa, Madagascar.
var. *dichroa*, Hort.
- S. Wallichii**, *Spring*. Tropical Asia.
- S. Warscewiczii**, Klotzsch. *See radiata*.
- S. watsoniana**, Hort. Garden origin.
- S. Willdenovii**, Baker. Tropical Asia. *Lycopodium Willdenovii*, Desv. ; Hook. & Grev. Ic. Fil. t. 57.

Tmesipteris, Bernh.

T. Fosteri, Endl. *See tannensis*.

T. tannensis, Bernh. Australia, New Zealand, Polynesia. *T. truncata*, Bernh.; Hook. Gen. Fil. t. 86.

III.—APPENDIX.

Cultural Forms of British Ferns.

Adiantum, L.

A. Capillus-Veneris, L.

- var. *daphnites*, *Lowe*.
- var. *elegantissimum*, *Hort.*
- var. *grande*, *Moore*.
- var. *imbricatum*, *Lowe*.
- var. *Mariesii*, *Moore*.
- var. *obrienianum*, *Birkenhead*.
- var. *semilunatum*, *Lowe*.

Aspidium, Sw.

A. aculeatum, Sw.

- var. *Braunii*, *Hort.*
- var. *cruciatum*, *Jones*.
- var. *densum*, *Moore*.
- var. *furcans*, *Hort.*
- var. *grandiceps*, *Jones*.
- var. *multifidum*, *Woll.*
- var. *proliferum*, *Woll.*
- var. *pulcherrimum*, *Jones*.
- var. *subcristatum*, *Jones*.

A. *angulare, Willd.

* This is a sub-species of *A. aculeatum*, but for garden purposes it is more convenient to treat it as a species.

Aspidium, Sw.—cont.**A. angulare, Willd.—cont.**

- var. **acutilobum**, *Woll.* Lowe, Nat. Ferns, i. 127.
- var. **acutilobum acutum**, *Lowe.*
- var. **acutilobum proliferum**, *Moore.*
- var. **acuto-dissectum**, *Carbonell.*
- var. **acuto-divaricatum**, *Carbonell.*
- var. **assimile**, *Jones.*
- var. **athyrioides**, *Woll.*
- var. **Bayliæ**, *Lyell.*
- var. **brachiatum**, *Moore.* Lowe, Nat. Ferns, i. 106.
- var. **brachiatum cristatum**, *Jarvis.*
- var. **Buckleæ**, *Lowe.*
- var. **calomelanos**, *Jones.*
- var. **capitatum**, *Clap.*
- var. **caudatum**, *Carbonell.*
- var. **Claphami**, *Jones.*
- var. **concinnum**, *Moore.* Lowe, Nat. Ferns, i. 173.
- var. **confluens**, *Moore.* Lowe, Nat. Ferns, i. 167.
- var. **congestum**, *Moore.* Lowe, Nat. Ferns, i. 120.
- var. **conspicuilebolum**, *Lowe.*
- var. **conspicuilebolum acutum**, *Carbonell.*
- var. **cristatum**, *Moore.* Lowe, Nat. Ferns, i. t. 16.
- var. **cristatum clintonianum**, *Hort.*
- var. **cristatum gracile**, *Woll.*
- var. **cristatum grandiceps**, *Hort.*
- var. **cruciato-polydactylum**, *Jones.*
- var. **cruciatum**, *Lowe.*
- var. **cruciatum proliferum**, *Hort.*
- var. **decompositum**, *Moore.* Lowe, Nat. Ferns, i. 79.
- var. **decompositum densum**, *Carbonell.*
- var. **decompositum grande**, *Jones.*
- var. **decompositum plumosum**, *Jones.*

Aspidium, Sw.—cont.**A. angulare, Willd.—cont.**

- var. *decompositum polydactylum*, *Jones*.
- var. *decompositum robustum*, *Jones*.
- var. *decompositum splendens*, *Jones*. Lowe, Nat. Ferns, i. 174.
- var. *decurrens*, *Moore*. Lowe, Nat. Ferns, i. 98.
- var. *densum*, *Moore*. Lowe, Nat. Ferns, i. 96.
- var. *divisilobum*, *Woll*.
- var. *divisilobum acutum*, *Jones*.
- var. *divisilobum cristatum*, *Ivery*.
- var. *divisilobum decorum*, *Jones*.
- var. *divisilobum densum*, *Carbonell*.
- var. *divisilobum foliosum*, *Jones*.
- var. *divisilobum lineare*, *Jones*.
- var. *divisilobum multifidum*, *Jones*.
- var. *divisilobum plumosum*, *Lowe*.
- var. *divisilobum proliferum*, *Woll*.
- var. *divisilobum reflexum*, *Jones*.
- var. *divisilobum robustum*, *Jones*.
- var. *divisilobum splendidissimum*, *Jones*.
- var. *dubium*, *Woll*. Lowe, Nat. Ferns, i. 86.
- var. *flexuosum*, *Woll*. Lowe, Nat. Ferns, i. 133.
- var. *flexuosum polydactylum*, *Jones*.
- var. *folioso-decompositum*, *Jones*.
- var. *foliosum*, *Woll*. Lowe, Nat. Ferns, i. 117.
- var. *foliosum multifidum*, *Jones*.
- var. *frondosum*, *Jones*.
- var. *frondosum decompositum*, *Jones*.
- var. *frondosum polydactylum*, *Jones*.
- var. *grandiceps*, *Moore*. Lowe, Nat. Ferns, i. 150.
- var. *grandidens*, *Jones*. Lowe, Nat. Ferns, i. 85.
- var. *inæquale variegatum*, *Woll*.

Aspidium, Sw.—cont.**A. angulare, Willd.—cont.**

- var. *inciso-acutum*, *Stansf.* Lowe, Nat. Ferns, i. 137.
 var. *Kitsoniae*, *Moore*. Lowe, Nat. Ferns, i. 124.
 var. *laciniatum*, *Woll.* Lowe, Nat. Ferns, i. 94.
 var. *laciniatum flexuosum*, *Hort.*
 var. *lanceolatum*, *Moore*.
 var. *latifolium*, *Woll.*
 var. *latifolium grandiceps*, *Jones*.
 var. *laxo-crispum*, *Jones*.
 var. *lineare*, *Moore*. Lowe, Nat. Ferns, i. 93.
 var. *lineare polydactylum*, *Jones*.
 var. *lineatum*, *Moly*.
 var. *Marshallii*, *Hort.*
 var. *multifidum densum*, *Jones*.
 var. *multilobum*, *Woll.* Lowe, Nat. Ferns, i. 121.
 var. *multilobum densum*, *Jones*.
 var. *multilobum multifidum*, *Jones*.
 var. *multilobum ovale*, *Woll.*
 var. *multilobum polydactylum*, *Jones*.
 var. *multilobum revolvens*, *Jones*.
 var. *oxyphyllum*, *Moore*. Lowe, Nat. Ferns, i. 116.
 var. *Pateyi*, *Moore*.
 var. *pendens*, *Jones*.
 var. *perserratum*, *Jones*.
 var. *plumosum*, *Woll.* Lowe, Nat. Ferns, i. 113.
 var. *plumosum elegantissimum*, *Jones*.
 var. *polydactylum*, *Moore*. Lowe, Nat. Ferns, i. 114.
 var. *polydactylum capitatum*, *Jones*.
 var. *polydactylum cruciatum*, *Lowe*.
 var. *polydactylum grande*, *Jones*.
 var. *polydactylum grandiceps*, *Jones*.
 var. *polydactylum laxum*, *Jones*.

Aspidium, Sw.—cont.**A. angulare, Willd.—cont.**

- var. *polydactylum multifidum*, *Jones*.
- var. *polydactylum robustum*, *Jones*.
- var. *proliferum*, *Moore*. Lowe, Nat. Ferns, i. 80.
- var. *proliferum densum*, *Carbonell*.
- var. *proliferum holeanum*, *Moore*.
- var. *pulcherrimum*, *Moore*.
- var. *pulchrum*, *Moore*.
- var. *raino-varians*, *Jones*.
- var. *remoto-decurrans*, *Woll*.
- var. *revolvens*, *Moore*.
- var. *rotundum polydactylon*, *Hort*.
- var. *sculptum*, *Carbonell*.
- var. *Stansfieldii*, *Lowe*.
- var. *stipatum*, *Woll*.
- var. *stipitatum*, *Moore*.
- var. *subpinnatum*, *Hort*.
- var. *subplumosum*, *Woll*.
- var. *Thompsoniæ*, *Lowe*. Nat. Ferns, i. 140.
- var. *torto-cristatum*, *Padley*.
- var. *tripinnatum*, *Moore*. Lowe, Nat. Ferns, i. 78.
- var. *tripinnatum decompositum*, *Lowe*. Nat. Ferns, i. 172.
- var. *tripinnatum elegans*, *Hort*.
- var. *tripinnatum polydactylum*, *Jones*.
- var. *varians*, *Woll*. Lowe, Nat. Ferns, i. 105.
- var. *venustum*, *Moore*.
- var. *Wollastoni majus*, *Hort*.

Asplenium, L.**A. Filix-fœmina, Bernh.**

- var. *abasiphyllum*, *Lowe*.

Asplenium, L.—*cont.*

A *Filix-fœmina*, *Bernh.*—*cont.*

- var. *adpresso-cristatum*, *Jones*.
- var. *Andersoni*, *Hort.*
- var. *angustatum*, *Lowe*.
- var. *apuæforme*, *Moore*. Lowe, Nat. Ferns, ii. 26.
- var. *axminsterense*, *Lowe*.
- var. *cancellatum cristatum*, *Jones*.
- var. *caudiculatum*, *Mapp.*
- var. *ceratophyllum*, *Mapp.*
- var. *clarissimum*, *Jones*.
- var. *congestum cristatum*, *Stansf.*
- var. *conoides cristatum*, *Stansf.*
- var. *coronatum*, *Lowe*. Nat. Ferns, ii. 35.
- var. *corymbiferum*, *Moore*.
- var. *Craigii*, *Moore*.
- var. *Craigii laciniatum*, *Jones*.
- var. *crispum coronatum*, *Jones*. Lowe, Nat. Ferns., ii. 36.
- var. *crispum cristatum*, *Carbonell*.
- var. *cristato-cristatum*, *Hort.*
- var. *cristatum*, *Woll.*
- var. *cristatum capitatum*, *Carbonell*.
- var. *cristatum densum*, *Carbonell*.
- var. *cristatum reflexum*, *Carbonell*.
- var. *cruciato-capitatum*, *Carbonell*.
- var. *cruciato-coronatum*, *Carbonell*.
- var. *cruciato-cristatum*, *Jones*.
- var. *cruciato-gracile*, *Jones*.
- var. *cruciato-grandiceps*, *Jones*.
- var. *cruciato-magnificum*, *Jones*.
- var. *cruciato-multifidum*, *Stansf.*
- var. *cruciatum*, *Jones*.

Asplenium, L.—*cont.*

A. *Filix-fœmina*, *Bernh.*—*cont.*

- var. *cruciatum corymbiferum*, *Hort.*
- var. *curtum*, *Woll.* Lowe, Nat. Ferns, ii. 32.
- var. *curtum cristatum*, *Stansf.*
- var. *daphnites*, *Glave.*
- var. *daphnites capitatum*, *Glave.*
- var. *defecto-furcellans*, *Hort.*
- var. *defecto-gracile*, *Mapp.*
- var. *diffidum*, *Jones.*
- var. *diffiso-multifidum*, *Moore.* Lowe, Nat. Ferns, ii. 44.
- var. *digitale capitatum*, *Jones.*
- var. *digitale purpureum*, *Jones.*
- var. *Elworthii*, *Moore.* Lowe, Nat. Ferns, ii. 46.
- var. *Fieldiæ*, *Moore.* Lowe, Nat. Ferns, ii. 50.
- var. *Fieldiæ minus*, *Jones.*
- var. *fissidens*, *Moore.* Lowe, Nat. Ferns, ii. 52.
- var. *flabelliceps*, *Carbonell.*
- var. *flexuosum*, *Woll.* Lowe, Nat. Ferns, ii. 49.
- var. *folioso-crispatum*, *Carbonell.*
- var. *folioso-cristatum*, *Woll.*
- var. *Frizelliæ*, *Moore.* Lowe, Nat. Ferns, ii. 53.
- var. *Frizelliæ cristatum*, *Jones.* Lowe, Nat. Ferns, ii. 142.
- var. *Frizelliæ multifidum*, *Barnes.*
- var. *Frizelliæ nanum*, *Ivery.* Lowe, Nat. Ferns, ii. 54.
- var. *furcans*, *Moore.*
- var. *germanicum*, *Hort.*
- var. *glomeratum*, *Moore.* Lowe, Nat. Ferns, ii. 56.
- var. *glomeratum grandiceps*, *Jones.*
- var. *gracile*, *Moore.*
- var. *grammicon*, *Lowe.*
- var. *grandiceps*, *Moore.*

Asplenium, L.—cont.

A. *Filix-fœmina*, *Bernh.*—cont.

- var. *grandiceps globosum*, *Sim.*
- var. *grandiceps minus*, *Carbonell.*
- var. *grandiceps pumilum*, *Sim.*
- var. *Howardæ*, *Moore.*
- var. *imbricatum*, *Carbonell*
- var. *inæquale*, *Woll.*
- var. *inæquale cristatum*, *Jones.*
- var. *interruptum*, *Moore.* Lowe, Nat. Ferns, ii. 62.
- var. *irregulare*, *Moore.* Lowe, Nat. Ferns, ii. 63.
- var. *kalothrix*, *Lowe.*
- var. *laceratum*, *Moore.* Lowe, Nat. Ferns, ii. 66.
- var. *laciniatum*, *Moore.* Lowe, Nat. Ferns, ii. t. 39A.
- var. *laciniatum caudatum*, *Hort.*
- var. *laciniatum elegans*, *Stansf.*
- var. *laciniatum furcans*, *Jones.*
- var. *laciniatum multifidum*, *Lowe.*
- var. *Lowæ*, *Lowe.*
- var. *Lowæ capitatum*, *Lowe.*
- var. *Medusæ ramocristatum*, *Hort.*
- var. *multifidum*, *Moore.* Lowe, Nat. Ferns, ii. t. 36.
- var. *multifidum laxum*, *Jones.*
- var. *nodosum*, *Jones.*
- var. *nodosum multifidum*, *Jones.*
- var. *nodosum nanum*, *Jones.*
- var. *nodosum ramulosum*, *Jones.*
- var. *nodosum truncatum*, *Jones.*
- var. *orbiculatum*, *Lowe.* Nat. Ferns, ii. 127.
- var. *pannosum*, *Moore.* Lowe, Nat. Ferns, ii. 78.
- var. *pannosum cristatum*, *Jones.*
- var. *Parsonsiæ*, *Moore.* Lowe, Nat. Ferns, ii. 79.
- var. *paucidentatum cristatum*, *Mapp.*

Asplenium, L.—*cont.*

A. *Filix-fœmina*, *Bernh.*—*cont.*

- var. *pectinatum*, *Moore*.
- var. *pectinatum Stableri*, *Mapp.*
- var. *percrustatum*, *Lowe*.
- var. *perfectum*, *Carbonell*.
- var. *phyllomanes*, *Mapp.*
- var. *plumosum*, *Moore*.
- var. *plumosum divaricatum*, *Moore*.
- var. *plumosum Edwardsi*, *Sim.*
- var. *plumosum elegans*, *Parsons*.
- var. *plumosum Hodgsonæ*, *Lowe*.
- var. *polydactylon*, *Moore*.
- var. *polydactylon cristatum*, *Stansf.*
- var. *polydactylon grandiceps*, *Jones*.
- var. *proteoides*, *Lowe*.
- var. *proteum*, *Clap.*
- var. *proteum cristatum*, *Jones*.
- var. *pulcherrimum*, *Jones*.
- var. *pulcherrimum cristatum*, *Jones*.
- var. *Pulleri*, *Moore*.
- var. *purpureum corymbiferum*, *Hort.*
- var. *rectangulare*, *Moore*.
- var. *reflexum*, *Woll.*
- var. *refractum*, *Moore*.
- var. *regale*, *Moore*.
- var. *rhæticum*, *Deakin*. Lowe, Nat. Ferns, ii. 128.
- var. *rhæticum reflexum*, *Mapp.*
- var. *rotundatum*, *Woll.*
- var. *rotundatum lunulatum*, *Carbonell*.
- var. *setigerum*, *Jones*.
- var. *Stableri*, *Lowe*.
- var. *Stansfieldii*, *Lowe*.

Asplenium, L.—cont.**A Filix-fœmina, Bernh.—cont.**


- var. *stipatum*, *Moore*.
- var. *stipatum capitatum*, *Carbonell*.
- var. *subcruciatum*, *Hort*.
- var. *todeoides*, *Stansf*.
- var. *tortile*, *Moore*.
- var. *torto-cristatum*, *Carbonell*.
- var. *torto-grandiceps*, *Carbonell*.
- var. *truncato-pinnulum*, *Jones*.
- var. *Tyermannii*, *Hort*.
- var. *unco-cristatum*, *Jones*.
- var. *unco-ramulosum*, *Stansf*.
- var. *variegatum*, *Woll*.
- var. *Vernoniæ*, *Jervis*. Lowe, Nat. Ferns, ii. 93.
- var. *Vernoniæ corymbiferum*, *Stansf*.
- var. *Vernoniæ crispum*, *Stansf*.
- var. *Vernoniæ cristatum*, *Mapp*.
- var. *Victoriæ*, *Lowe*. Nat. Ferns, ii. 97.
- var. *Williamsii*, *Jones*.

Cystopteris, Bernh.**C. fragilis, Bernh.**

- var. *angustata*, *Smith*.
- var. *dickieana*, *Moore*.

Nephrodium, L. C. Rich.**N. Filix-mas, L. C. Rich.**

- var. *attenuatum*, *Padley*.
- var. *Barnesii*, *Moore*. Lowe, Nat. Ferns, i. 272.
- var. *Bollandæ*, *Moore*.
- var. *capitatum*, *Carbonell*.



*Nephrodium, L. C. Rich.—cont.**N. Filix-mas, L. C. Rich.—cont.*

- var. *concinnum*, *Hort.*
- var. *crispatissimum*, *Jones.*
- var. *crispulatum*, *Hort.*
- var. *crispum*, *Sim.* Lowe, Nat. Ferns, i. 248.
- var. *crispum gracile*, *Lyell.*
- var. *Crista-Galli*, *Carbonell.*
- var. *cristatum*, *Lowe.*
- var. *cristatum angustatum*, *Moore.*
- var. *cristatum caudatum*, *Ivery.*
- var. *cronkleyense*, *Lowe.*
- var. *depauperatum*, *Padley.*
- var. *digitatum*, *Lowe.*
- var. *Ellacombei*, *Lowe.*
- var. *erosum*, *Clowes.* Lowe, Nat. Ferns, i. 248.
- var. *flexuosum polydactylon*, *Hort.*
- var. *fluctuosum*, *Moore.*
- var. *furcans*, *Moore.* Lowe, Nat. Ferns, i. 244.
- var. *grandiceps*, *Sim.* Lowe, Nat. Ferns, i. 275.
- var. *Ingrami*, *Moore.*
- var. *interruptum*, *Moore.* Lowe, Nat. Ferns, i. 242.
- var. *lineare*, *Woll.*
- var. *multiforme*, *Barnes.*
- var. *paleaceum*, *Hook.*
- var. *paleaceum cristatum*, *Moore.*
- var. *paleaceum cristatum intermedium*, *Sim.*
- var. *paleaceum ramo-furcillatum*, *Woll.*
- var. *paleaceum revolvens*, *Woll.*
- var. *polydactylum*, *Lowe.* Nat. Ferns, i. 257.
- var. *propinquum*, *Woll.*
- var. *propinquum cristatum*, *Moore.*
- var. *propinquum gracile*, *Moore.*

Nephrodium, *L. C. Rich.*—*cont.*

N. Filix-mas, *L. C. Rich.*—*cont.*

- var. *propinquum grandiceps*, *Moore*.
- var. *propinquum imbricatum*, *Hort.*
- var. *propinquum parviceps*, *Barnes*.
- var. *propinquum polydactylum*, *Barnes*.
- var. *pseudo-mas*, *Woll.* See var. *paleaceum*.
- var. *Pulleri*, *Jones*.
- var. *ramo-cristatum*, *Stansf.*
- var. *ramo-furcillatum*, *Carbonell*.
- var. *ramosum*, *Barnes*.
- var. *revolvens*, *Humphreys*.
- var. *serratum cristatum*, *Hort.*
- var. *subcristatum*, *Carbonell*.
- var. *variegatum*, *Lowe*.
- var. *willyanum*, *Hort.*

N. spinulosum, *Desv.*

- var. *dilatatum*, *Hook. Brit. Ferns*, t. 19.
- var. *dilatatum angusti-pinnulum*, *Sim. Lowe*, *Nat. Ferns*, i. 313.
- var. *dilatatum calomelanos*, *Jones*.
- var. *dilatatum corymbiferum*, *Hort.*
- var. *dilatatum foliosum multicristatum*, *Hort.*
- var. *dilatatum foliosum ramosum*, *Hort.*
- var. *dilatatum grandiceps*, *Barnes*.
- var. *dilatatum Howardæ*, *Jones*.
- var. *dilatatum lepidotum*, *Moore*.

Osmunda, *L.*

O. regalis, *L.*

- var. *cristata*, *Moore*.
- var. *palustris*, *Hort.*
- var. *purpurascens*, *Moore*.



Polypodium, L.**P. Phegopteris, L.**var. *cristatum*, *Jones*.var. *strictum*, *Barnes*.**P. vulgare, L.**var. *acuto-serratum*, *Jones*.var. *acutum*, *Moore*. Lowe, Nat. Ferns, i. 27.var. *cambricum*, *Willd*. Lowe, Nat. Ferns, i. 32.var. *cambricum Barrowi*, *Barnes*.var. *cornubiense*, *Moore*.var. *cornubiense elegantissimum*, *Stansf*.var. *Elworthii*, *Ivery*.var. *falcatum*, *Jones*.var. *grandiceps*, *Hort*.var. *macrostachya*, *O'Kelly*.var. *multifido-cristatum*, *Stansf*. Lowe, Nat. Ferns, i. 43.var. *omnilacerum*, *Moore*. Lowe, Nat. Ferns, i. t. 7B.var. *pulcherrimum*, *Moore*. Lowe, Nat. Ferns, i. 46.var. *ramosum*, *Moore*.var. *semilacerum*, *Link*. Lowe, Nat. Ferns, i. 47.var. *serratum*, *Moore*. Lowe, Nat. Ferns, i. 48.var. *serratum elegans*, *Hort*.var. *trichomanoides*, *Backh*.**Scolopendrium, J. E. Sm.****S. vulgare, J. E. Sm.**var. *aduncum*, *Moore*. Lowe, Nat. Ferns, ii. 354.var. *crispum*, *Grey*.var. *Crista-Galli*, *Woll*.var. *Cuspis*, *Carbonell*.var. *fisso-crenatum*, *Jones*.

Scolopendrium, J. E. Sm.—cont.

S. vulgare, J. E. Sm.—cont.

var. *grandiceps*, *Jones*.

var. *illustre*, *Lowe*.

var. *involvens*, *Mapp.*

var. *labyrinthicum*, *Lowe*.

var. *laceratum*, *Moore*.

var. *macrosorum*, *Moore*.

var. *marginatum*, *Moore*.

var. *Œnone*, *Hort.*

var. *peraferens*, *Woll.*

var. *peraferens ovale*, *Hort*

var. *ramulosissimum*, *Woll.*

var. *sculpturatum*, *Moore*. Lowe, Nat. Ferns, ii. 249.

var. *submarginato-fissum*, *Wills*.

var. *undulatum cochleatum*, *Jones*.

var. *variegatum*, *Woll.* Lowe, Nat. Ferns, ii. 333.

var. *Veloisii*, *Moore*.

ROYAL BOTANIC GARDENS, KEW.

H A N D - L I S T

FERNS AND FERN ALLIES

CULTIVATED IN

THE ROYAL BOTANIC GARDENS.

(SECOND EDITION.)




L O N D O N :


SOLD AT THE ROYAL BOTANIC GARDENS, KEW.
PRINTED FOR HIS MAJESTY'S STATIONERY OFFICE.
BY DARLING & SON, LTD., 34-40, BACON STREET, E.

—
1906.

Price Fivepence.







QK Kew, Eng. Royal Botanic
73 Gardens
K4A237 Hand-list of ferns
1906

BioMed

PLEASE DO NOT REMOVE
CARDS OR SLIPS FROM THIS POCKET

UNIVERSITY OF TORONTO LIBRARY
